



PRIOR AUTHORIZATION POLICY

- POLICY:** Immunologicals – Dupixent Prior Authorization Policy
- Dupixent® (dupilumab subcutaneous injection – Regeneron/sanofi-aventis)

REVIEW DATE: 04/19/2024; selected revisions 09/25/2024 and 10/09/2024

INSTRUCTIONS FOR USE

THE FOLLOWING COVERAGE POLICY APPLIES TO HEALTH BENEFIT PLANS ADMINISTERED BY CIGNA COMPANIES. CERTAIN CIGNA COMPANIES AND/OR LINES OF BUSINESS ONLY PROVIDE UTILIZATION REVIEW SERVICES TO CLIENTS AND DO NOT MAKE COVERAGE DETERMINATIONS. REFERENCES TO STANDARD BENEFIT PLAN LANGUAGE AND COVERAGE DETERMINATIONS DO NOT APPLY TO THOSE CLIENTS. COVERAGE POLICIES ARE INTENDED TO PROVIDE GUIDANCE IN INTERPRETING CERTAIN STANDARD BENEFIT PLANS ADMINISTERED BY CIGNA COMPANIES. PLEASE NOTE, THE TERMS OF A CUSTOMER'S PARTICULAR BENEFIT PLAN DOCUMENT [GROUP SERVICE AGREEMENT, EVIDENCE OF COVERAGE, CERTIFICATE OF COVERAGE, SUMMARY PLAN DESCRIPTION (SPD) OR SIMILAR PLAN DOCUMENT] MAY DIFFER SIGNIFICANTLY FROM THE STANDARD BENEFIT PLANS UPON WHICH THESE COVERAGE POLICIES ARE BASED. FOR EXAMPLE, A CUSTOMER'S BENEFIT PLAN DOCUMENT MAY CONTAIN A SPECIFIC EXCLUSION RELATED TO A TOPIC ADDRESSED IN A COVERAGE POLICY. IN THE EVENT OF A CONFLICT, A CUSTOMER'S BENEFIT PLAN DOCUMENT ALWAYS SUPERSEDES THE INFORMATION IN THE COVERAGE POLICIES. IN THE ABSENCE OF A CONTROLLING FEDERAL OR STATE COVERAGE MANDATE, BENEFITS ARE ULTIMATELY DETERMINED BY THE TERMS OF THE APPLICABLE BENEFIT PLAN DOCUMENT. COVERAGE DETERMINATIONS IN EACH SPECIFIC INSTANCE REQUIRE CONSIDERATION OF 1) THE TERMS OF THE APPLICABLE BENEFIT PLAN DOCUMENT IN EFFECT ON THE DATE OF SERVICE; 2) ANY APPLICABLE LAWS/REGULATIONS; 3) ANY RELEVANT COLLATERAL SOURCE MATERIALS INCLUDING COVERAGE POLICIES AND; 4) THE SPECIFIC FACTS OF THE PARTICULAR SITUATION. EACH COVERAGE REQUEST SHOULD BE REVIEWED ON ITS OWN MERITS. MEDICAL DIRECTORS ARE EXPECTED TO EXERCISE CLINICAL JUDGMENT AND HAVE DISCRETION IN MAKING INDIVIDUAL COVERAGE DETERMINATIONS. COVERAGE POLICIES RELATE EXCLUSIVELY TO THE ADMINISTRATION OF HEALTH BENEFIT PLANS. COVERAGE POLICIES ARE NOT RECOMMENDATIONS FOR TREATMENT AND SHOULD NEVER BE USED AS TREATMENT GUIDELINES. IN CERTAIN MARKETS, DELEGATED VENDOR GUIDELINES MAY BE USED TO SUPPORT MEDICAL NECESSITY AND OTHER COVERAGE DETERMINATIONS.

CIGNA NATIONAL FORMULARY COVERAGE:

OVERVIEW

Dupixent, an interleukin-4 receptor alpha antagonist, is indicated for the following uses:¹

- **Asthma**, as an add-on maintenance treatment in patients ≥ 6 years of age with moderate-to-severe disease with an eosinophilic phenotype or with oral corticosteroid-dependent asthma.
Limitation of Use: Dupixent is not indicated for the relief of acute bronchospasm or status asthmaticus.
- **Atopic dermatitis**, for the treatment of patients ≥ 6 months of age with moderate-to-severe disease not adequately controlled with topical prescription therapies or when those therapies are not advisable.
- **Chronic obstructive pulmonary disease (COPD)**, as add-on maintenance treatment in patients ≥ 18 years of age with inadequately controlled disease and an eosinophilic phenotype.
Limitation of Use: Dupixent is not indicated for the relief of acute bronchospasm.
- **Chronic rhinosinusitis with nasal polyposis (CRSwNP)** [i.e., nasal polyps], as an add-on maintenance treatment in patients ≥ 12 years of age with inadequately controlled disease.
- **Eosinophilic esophagitis**, in patients ≥ 1 year of age who weigh ≥ 15 kg.
- **Prurigo nodularis**, in patients ≥ 18 years of age.

Clinical Efficacy

Asthma

Timing of efficacy assessments varied by indication across the numerous pivotal studies in which Dupixent demonstrated benefit. In the asthma trials, efficacy with Dupixent was assessed as early as 24 weeks.²⁻⁵

Atopic Dermatitis

In atopic dermatitis, the majority of studies evaluated the efficacy of Dupixent at 16 weeks.^{1,6-10}

Chronic Obstructive Pulmonary Disease

Two pivotal studies evaluated Dupixent in adults with COPD.^{32,33} To be eligible for enrollment, patients had a blood eosinophil level ≥ 300 cells per microliter. Patients were required to have been receiving background triple inhaler therapy (i.e., an inhaled corticosteroid [ICS] with a long-acting muscarinic antagonist [LAMA] and a long-acting beta2-agonist [LABA]) or LAMA/LABA combination therapy if the patient had an ICS contraindication, for at least 3 months prior to randomization. Patients also had signs or symptoms of chronic bronchitis for ≥ 3 months and had experienced at least two moderate COPD exacerbations or one severe COPD exacerbation the year prior to screening. Moderate COPD exacerbations were defined as exacerbations that required treatment with a systemic corticosteroid, an antibiotic agent, or both. At least one of the moderate exacerbations had to have required treatment with a systemic corticosteroid. Severe exacerbations were defined as exacerbations that led to hospitalization or an emergency medical care visit with observation lasting > 24 hours. Overall, at least one of the patient's exacerbations had to have occurred while they were receiving ICS/LAMA/LABA therapy (or a LAMA/LABA if the patient had an ICS contraindication). Patients were randomized to receive either Dupixent or placebo in addition to background maintenance therapy (i.e., ICS/LAMA/LABA triple therapy or LAMA/LABA therapy if the patient had an ICS contraindication) for 52 weeks. While lung function parameters were improved as early as Week 12 (3 months), the other major efficacy endpoints were evaluated at Week 52 (e.g., exacerbations, dyspnea scores).

Chronic Rhinosinusitis with Nasal Polyps

The pivotal studies involving patients with CRSwNP evaluated the primary efficacy endpoints following 24 weeks of treatment.^{1,11-13} Patients continued treatment with intranasal corticosteroids throughout the studies.

Eosinophilic Esophagitis

In Dupixent's eosinophilic esophagitis pivotal study, patients ≥ 12 years of age were required to have disease confirmed by baseline endoscopic biopsies with a demonstration of eosinophilic infiltration on central reading (peak cell count ≥ 15 eosinophils per high-powered field) that was unresponsive to an 8 week course of treatment with a high-dose proton pump inhibitor.¹⁴ Patients with other causes of eosinophilic esophagitis, such as hypereosinophilic syndrome and eosinophilic granulomatosis with polyangiitis, were excluded from the study. In the first portion of this study, efficacy, as measured by objective assessments (e.g., intraepithelial eosinophil count) and subjective assessments (e.g., dysphagia symptoms), was evaluated after 24 weeks (6 months) of Dupixent therapy. A very similarly designed pivotal study evaluated the efficacy of Dupixent for the treatment of eosinophilic esophagitis in patients 1 to 11 years of age.¹ Endoscopic biopsy evidence of eosinophilic infiltration despite treatment with a proton pump inhibitor was again required for study enrollment.

Prurigo Nodularis

Two pivotal studies, PRIME and PRIME2, evaluated Dupixent's efficacy in the treatment of prurigo nodularis.^{15,16} To enroll, patients were required to have ≥ 20 identifiable nodular

lesions in total on both legs, and/or both arms, and/or trunk and to have failed a 2-week trial of a topical corticosteroid. Patients with prurigo nodularis secondary to medications or a medical condition such as neuropathy or psychiatric disease were excluded from the studies. The primary endpoint was evaluated at Week 24 in PRIME and initially at Week 12 and again at Week 24 in PRIME2.

Guidelines

Asthma Guidelines

The Global Initiative for Asthma Global Strategy for Asthma Management and Prevention (2023) proposes a stepwise approach to asthma treatment.¹⁷ Dupixent is listed as an option for add-on therapy in patients ≥ 6 years of age with severe eosinophilic/Type 2 asthma or for patients ≥ 12 years of age who require treatment with a maintenance oral corticosteroid. Higher blood eosinophil levels and higher fractional concentration of exhaled nitric oxide may predict a good asthma response to Dupixent.

According to the European Respiratory Society/American Thoracic Society guidelines (2014; updated in 2020), severe asthma is defined as asthma which requires treatment with a high-dose inhaled corticosteroid (ICS) in addition to a second controller medication (and/or systemic corticosteroids) to prevent it from becoming uncontrolled, or asthma which remains uncontrolled despite this therapy.^{18,19} Uncontrolled asthma is defined as asthma that worsens upon tapering of high-dose ICS or systemic corticosteroids or asthma that meets one of the following four criteria:

- 1) Poor symptom control: Asthma Control Questionnaire consistently ≥ 1.5 or Asthma Control Test < 20 ; OR
- 2) Frequent severe exacerbations: two or more bursts of systemic corticosteroids in the previous year; OR
- 3) Serious exacerbations: at least one hospitalization, intensive care unit stay, or mechanical ventilation in the previous year; OR
- 4) Airflow limitation: forced expiratory volume in 1 second (FEV_1) $< 80\%$ predicted after appropriate bronchodilator withholding.

Atopic Dermatitis Guidelines

Guidelines for the care and management of atopic dermatitis (with topical therapies in adults [2022], with phototherapy and systemic agents [2023]) have been updated to address Dupixent.^{20,21} The guidelines note that despite the availability of newer, systemic therapies (e.g., Dupixent), topical agents remain the mainstay of treatment due to their proven track record and favorable safety profiles. Several topical agents are recommended, with topical corticosteroids commonly used first-line for mild to severe atopic dermatitis in all skin regions. If topical therapy and basic management (e.g., moisturizers, bathing modifications) have been optimized and the patient has not achieved adequate control, consider an alternative diagnosis or systemic therapy. In this setting, use of Dupixent is recommended in patients with moderate to severe disease (strong recommendation).

Chronic Obstructive Pulmonary Disease Guidelines

The Global Initiative for Chronic Obstructive Lung Disease (GOLD) Global Strategy for the Diagnosis, Management, and Prevention of Chronic Obstructive Pulmonary Disease (2024) does not address Dupixent.³⁴ Triple therapy with an ICS/LAMA/LABA combination is recommended in patients with a history of exacerbations and elevated eosinophils. However, guidelines note that ICS therapy increases the risk of pneumonia in patients with COPD, particularly those with severe disease.

Chronic Rhinosinusitis with Nasal Polyps Guidelines

The Joint Task Force on Practice Parameters (JTFFP) published a focused guideline update for the medical management of CRSwNP (2023), which updated recommendations regarding intranasal corticosteroids and biologic therapies.²² Intranasal corticosteroids are recommended for the treatment of CRSwNP. Use of biologics (e.g., Dupixent) is also recommended. However, in patients who derived a sufficient benefit from other therapies such as intranasal corticosteroids, surgery, or aspirin therapy after desensitization, biologics may not be preferred. Conversely, biologics may be preferred over other medical treatment options in patients who continue to have a high burden of disease despite receiving at least 4 weeks of treatment with an intranasal corticosteroid.

The diagnosis of CRSwNP was not addressed in this focused guideline update, but previous guidelines have noted that the presence of two or more signs and symptoms of chronic rhinosinusitis (e.g., rhinorrhea, postnasal drainage, anosmia, nasal congestion, facial pain, headache, fever, cough, and purulent discharge) that persist for an extended period of time makes the diagnosis chronic rhinosinusitis likely.²³⁻²⁶ However, this requires confirmation of sinonasal inflammation, which can either be done via direct visualization or computed tomography (CT) scan. Oral corticosteroids and surgical intervention were not specifically addressed in this update, but prior guidelines recommend short courses of oral corticosteroid as needed and consideration of surgical removal as an adjunct to medical therapy in patients with CRSwNP that is not responsive or is poorly responsive to medical therapy.^{23,24,26}

Eosinophilic Esophagitis Guidelines

Guidelines for the management of eosinophilic esophagitis from the American Gastroenterological Association and the Joint Task Force on Allergy-Immunology Practice Parameters (2020) have not been updated since the FDA approval of Dupixent for this indication.²⁷ In patients with symptomatic disease, use of a proton pump inhibitor is recommended over no treatment, as is treatment with topical corticosteroids. Guidelines recommend diet modifications, such as an elemental diet (amino acid based formulas) or an elimination diet, over no treatment. However, it is noted that patients who put a higher value on avoiding the challenges of adherence to these diets and the prolonged process of dietary reintroduction may reasonably decline this treatment option.

Prurigo Nodularis Guidelines

A United States Expert Panel Consensus provides a practical approach for the diagnosis and management of prurigo nodularis (2021).²⁸ The primary findings in patients with prurigo nodularis are the presence of firm, nodular lesions; pruritus lasting at least 6 weeks; and history or signs, or both, of repeated scratching, picking, or rubbing. Goals of treatment are to reduce pruritus, interrupt the itch-scratch cycle, and completely heal prurigo nodularis lesions.

POLICY STATEMENT

Prior Authorization is recommended for prescription benefit coverage of Dupixent. All approvals are provided for the duration noted below. In cases where the approval is authorized in months, 1 month is equal to 30 days. Because of the specialized skills required for evaluation and diagnosis of patients treated with Dupixent as well as the monitoring required for adverse events and long-term efficacy, initial approval requires Dupixent to be prescribed by or in consultation with a physician who specializes in the condition being treated.

- **Dupixent® (dupilumab subcutaneous injection – Regeneron/sanofi-aventis)**

is(are) covered as medically necessary when the following criteria is(are) met for FDA-approved indication(s) or other uses with supportive evidence (if applicable):

FDA-Approved Indications

- 1. Asthma.** Approve for the duration noted if the patient meets ONE of the following (A or B):
 - A) Initial Therapy.** Approve for 6 months if the patient meets ALL of the following (i, ii, iii, iv, and v):
 - i.** Patient is ≥ 6 years of age; AND
 - ii.** Patient meets ONE of the following (a or b):
 - a) Patient meets ONE of the following (1 or 2):**
 - (1)** Patient has a blood eosinophil level ≥ 150 cells per microliter within the previous 6 weeks; OR
 - (2)** Patient had a blood eosinophil level ≥ 150 cells per microliter prior to treatment with Dupixent or another monoclonal antibody therapy that may alter blood eosinophil levels; OR

Note: Examples of monoclonal antibody therapies that may alter blood eosinophil levels include Dupixent, Adbry (tralokinumab-ldrm subcutaneous injection), Cinqair (reslizumab intravenous infusion), Ebglyss (lebrikizumab-lbkz subcutaneous injection), Fasentra (benralizumab subcutaneous injection), Nemluvio (nemolizumab-ilto subcutaneous injection), Nucala (mepolizumab subcutaneous injection), Tezspire (tezepelumab subcutaneous injection), and Xolair (omalizumab subcutaneous injection).
 - b) According to the prescriber, the patient has oral (systemic) corticosteroid-dependent asthma (e.g., the patient has received ≥ 5 mg oral prednisone or equivalent per day for ≥ 6 months); AND**
 - iii.** Patient has received at least 3 consecutive months of combination therapy with BOTH of the following (a and b):
 - a) An inhaled corticosteroid; AND**
 - b) At least one additional asthma controller or asthma maintenance medication; AND**

Note: Examples of additional asthma controller or asthma maintenance medications are inhaled long-acting beta₂-agonists, inhaled long-acting muscarinic antagonists, and monoclonal antibody therapies for asthma (e.g., Cinqair, Fasentra, Nucala, Tezspire, and Xolair). Use of a combination inhaler containing both an inhaled corticosteroid and additional asthma controller/maintenance medication(s) would fulfill the requirement for both criteria a and b.
 - iv.** Patient has asthma that is uncontrolled or was uncontrolled at baseline as defined by ONE of the following (a, b, c, d, or e):

Note: "Baseline" is defined as prior to receiving Dupixent or another monoclonal antibody therapy for asthma. Examples of monoclonal antibody therapies for asthma include Dupixent, Cinqair, Fasentra, Nucala, Tezspire, and Xolair.

 - a) Patient experienced two or more asthma exacerbations requiring treatment with systemic corticosteroids in the previous year; OR**

- b) Patient experienced one or more asthma exacerbation(s) requiring a hospitalization, an emergency department visit, or an urgent care visit in the previous year; OR
 - c) Patient has a forced expiratory volume in 1 second (FEV₁) < 80% predicted; OR
 - d) Patient has an FEV₁/forced vital capacity (FVC) < 0.80; OR
 - e) Patient has asthma that worsens upon tapering of oral (systemic) corticosteroid therapy; AND
 - v. The medication is prescribed by or in consultation with an allergist, immunologist, or pulmonologist.
- B) Patient is Currently Receiving Dupixent.** Approve for 1 year if the patient meets ALL of the following (i, ii, and iii):
- i. Patient has already received at least 6 months of therapy with Dupixent; AND
Note: A patient who has received < 6 months of therapy or who is restarting therapy with Dupixent should be considered under criterion 1A (Asthma, Initial Therapy).
 - ii. Patient continues to receive therapy with one inhaled corticosteroid or one inhaled corticosteroid-containing combination inhaler; AND
 - iii. Patient has responded to therapy as determined by the prescriber.
Note: Examples of a response to Dupixent therapy are decreased asthma exacerbations; decreased asthma symptoms; decreased hospitalizations, emergency department visits, or urgent care visits due to asthma; decreased requirement for oral corticosteroid therapy.
- 2. Atopic Dermatitis.** Approve for the duration noted if the patient meets ONE of the following (A or B):
- A) Initial Therapy.** Approve for 4 months if the patient meets ALL of the following (i, ii, iii, and iv):
- i. Patient is ≥ 6 months of age; AND
 - ii. According to the prescriber, the patient has atopic dermatitis involvement estimated to be ≥ 10% of the body surface area; AND
 - iii. Patient meets ALL of the following (a, b, and c):
 - a) Patient has tried at least one medium-, medium-high, high-, and/or super-high-potency prescription topical corticosteroid; AND
 - b) This topical corticosteroid was applied daily for at least 28 consecutive days; AND
 - c) According to the prescriber, inadequate efficacy was demonstrated with this topical corticosteroid therapy; AND
 - iv. The medication is prescribed by or in consultation with an allergist, immunologist, or dermatologist.
- B) Patient is Currently Receiving Dupixent.** Approve for 1 year if the patient meets BOTH of the following (i and ii):
- i. Patient has already received at least 4 months of therapy with Dupixent; AND
Note: A patient who has received < 4 months of therapy or who is restarting therapy with Dupixent should be considered under criterion 2A (Atopic Dermatitis, Initial Therapy).
 - ii. Patient has responded to therapy as determined by the prescriber.
Note: Examples of a response to Dupixent therapy are marked improvements in erythema, induration/papulation/edema, excoriations, and lichenification; reduced pruritus; decreased requirement for other topical or systemic therapies; reduced body surface area affected with atopic dermatitis; or other responses observed.

3. Chronic Obstructive Pulmonary Disease (COPD). Approve for the duration noted if the patient meets ONE of the following (A or B):

A) Initial Therapy. Approve for 6 months if the patient meets ALL of the following (i, ii, iii, iv, v, and vi):

i. Patient is ≥ 18 years of age; AND

ii. Patient meets ONE of the following (a or b):

a) Patient has a blood eosinophil level ≥ 300 cells per microliter within the previous 6 weeks; OR

b) Patient had a blood eosinophil level ≥ 300 cells per microliter prior to treatment with Dupixent or another monoclonal antibody therapy that may alter blood eosinophil levels; AND

Note: Examples of monoclonal antibody therapies that may alter blood eosinophil levels include Dupixent, Adbry (tralokinumab-ldrm subcutaneous injection), Cinqair (reslizumab intravenous infusion), Ebglyss (lebrikizumab-lbkz subcutaneous injection); Fasentra (benralizumab subcutaneous injection), Nemluvio (nemolizumab-ilto subcutaneous injection); Nucala (mepolizumab subcutaneous injection), Tezspire (tezepelumab subcutaneous injection), and Xolair (omalizumab subcutaneous injection).

iii. Patient meets ONE of the following (a or b):

a) Patient has received at least 3 consecutive months of combination therapy with ALL of the following (1, 2, and 3):

(1)Inhaled long-acting beta₂-agonist (LABA); AND

(2)Inhaled long-acting muscarinic antagonist (LAMA); AND

(3)Inhaled corticosteroid (ICS); OR

Note: Use of single-entity inhalers or a combination inhaler containing multiple agents from the medication classes listed would fulfill the requirement.

b) Patient meets BOTH of the following (1 and 2):

(1)Patient has received at least 3 consecutive months of combination therapy with an inhaled LABA and an inhaled LAMA; AND

Note: Use of single-entity inhalers or a combination inhaler containing multiple agents from the medication classes listed would fulfill the requirement.

(2)According to the prescriber, the patient has a contraindication to the use of an inhaled corticosteroid; AND

iv. According to the prescriber, the patient has had signs or symptoms of chronic bronchitis (e.g., chronic productive cough) for ≥ 3 months in the previous 12 months; AND

v. Patient meets ONE of the following (a or b):

a) Patient meets ALL of the following (1, 2, and 3):

(1)Patient experienced two or more COPD exacerbations requiring treatment with a systemic corticosteroid and/or an antibiotic in the previous 12 months; AND

(2)One or more of these COPD exacerbations required treatment with a systemic corticosteroid; AND

(3)One or more of these COPD exacerbations occurred while the patient was receiving combination therapy with an ICS, LAMA, and LABA or with a LAMA and LABA, if the patient has a contraindication to an ICS; OR

b) Patient meets ALL of the following (1 and 2):

(1)Patient experienced one or more COPD exacerbation(s) requiring a hospitalization in the previous 12 months; AND

Note: A hospitalization includes a hospital admission or an emergency medical care visit with observation lasting > 24 hours.

- (2) One or more of these COPD exacerbations occurred while the patient was receiving combination therapy with an ICS, LAMA, and LABA or with a LAMA and LABA, if the patient has a contraindication to an ICS; AND
- vi. The medication is prescribed by or in consultation with an allergist, immunologist, or pulmonologist.
- B) Patient is Currently Receiving Dupixent.** Approve for 1 year if the patient meets the following (i, ii, and iii):
- i. Patient has already received at least 6 months of therapy with Dupixent; AND Note: A patient who has received < 6 months of therapy or who is restarting therapy with Dupixent should be considered under criterion 3A (Chronic Obstructive Pulmonary Disease, Initial Therapy).
 - ii. Patient continues to receive combination therapy with an inhaled LABA and LAMA; AND Note: Use of single-entity inhalers or a combination inhaler containing multiple agents from the medication classes listed would fulfill the requirement.
 - iii. Patient has experienced a beneficial clinical response, defined by ONE of the following (a, b, c, d, or e):
 - a) Reduced COPD symptoms; OR
 - b) Reduced COPD exacerbations; OR
 - c) Reduced COPD-related hospitalizations; OR
 - d) Reduced emergency department or urgent care visits; OR
 - e) Improved lung function parameters.
- 4. Chronic Rhinosinusitis with Nasal Polyps.** Approve for the duration noted if the patient meets ONE of the following (A or B):
- A) Initial Therapy.** Approve for 6 months if the patient meets ALL of the following (i, ii, iii, iv, v, and vi):
- i. Patient is \geq 12 years of age; AND
 - ii. Patient has chronic rhinosinusitis with nasal polyps as evidenced by direct examination, endoscopy, or sinus computed tomography (CT) scan; AND
 - iii. Patient has experienced two or more of the following symptoms for at least 6 months: nasal congestion, nasal obstruction, nasal discharge, and/or reduction/loss of smell; AND
 - iv. Patient meets BOTH of the following (a and b):
 - a) Patient has received at least 4 weeks of therapy with an intranasal corticosteroid; AND
 - b) Patient will continue to receive therapy with an intranasal corticosteroid concomitantly with Dupixent; AND
 - v. Patient meets ONE of the following (a, b, or c):
 - a) Patient has received at least one course of treatment with a systemic corticosteroid for 5 days or more within the previous 2 years; OR
 - b) Patient has a contraindication to systemic corticosteroid therapy; OR
 - c) Patient has had prior surgery for nasal polyps; AND
 - vi. The medication is prescribed by or in consultation with an allergist, immunologist, or an otolaryngologist (ear, nose, and throat [ENT] physician specialist).
- B) Patient is Currently Receiving Dupixent.** Approve for 1 year if the patient meets ALL of the following (i, ii, and iii):
- i. Patient has already received at least 6 months of therapy with Dupixent; AND Note: A patient who has received < 6 months of therapy or who is restarting therapy with Dupixent should be considered under criterion 4A (Chronic Rhinosinusitis with Nasal Polyps, Initial Therapy).
 - ii. Patient continues to receive therapy with an intranasal corticosteroid; AND
 - iii. Patient has responded to therapy as determined by the prescriber.

Note: Examples of a response to Dupixent therapy are reduced nasal polyp size, improved nasal congestion, reduced sinus opacification, decreased sinonasal symptoms, improved sense of smell.

5. Eosinophilic Esophagitis. Approve for the duration noted if the patient meets ONE of the following (A or B):

A) Initial Therapy. Approve for 6 months if the patient meets ALL of the following (i, ii, iii, iv, v, vi and vii):

- i.** Patient is \geq 1 year of age; AND
- ii.** Patient weighs \geq 15 kg; AND
- iii.** Patient has a diagnosis of eosinophilic esophagitis as confirmed by an endoscopic biopsy demonstrating \geq 15 intraepithelial eosinophils per high-power field; AND
- iv.** Patient does not have a secondary cause of eosinophilic esophagitis; AND
Note: Examples of secondary causes of eosinophilic esophagitis are hypereosinophilic syndrome, eosinophilic granulomatosis with polyangiitis, and food allergy.
- v.** Patient has received at least 8 weeks of therapy with a proton pump inhibitor; AND
- vi.** Patient meets ONE of the following (a or b):
 - a)** Patient has tried dietary modifications to treat/manage eosinophilic esophagitis; OR
 - b)** The provider has determined that the patient is not an appropriate candidate for dietary modifications; AND
Note: Examples of dietary modifications to treat eosinophilic esophagitis include an elemental diet or an elimination diet.
- vii.** The medication is prescribed by or in consultation with an allergist or gastroenterologist.

B) Patient is Currently Receiving Dupixent. Approve for 1 year if the patient meets BOTH of the following (i and ii):

- i.** Patient has already received at least 6 months of therapy with Dupixent; AND
Note: A patient who has received $<$ 6 months of therapy or who is restarting therapy with Dupixent should be considered under criterion 5A (Eosinophilic Esophagitis, Initial Therapy).
- ii.** Patient has experienced a beneficial clinical response, defined by ONE of the following (a, b, or c):
 - a)** Reduced intraepithelial eosinophil count; OR
 - b)** Decreased dysphagia/pain upon swallowing; OR
 - c)** Reduced frequency/severity of food impaction.

6. Prurigo Nodularis. Approve for the duration noted if the patient meets ONE of the following (A or B):

A) Initial Therapy. Approve for 6 months if the patient meets ALL of the following (i, ii, iii, iv, v, and vi):

- i.** Patient is \geq 18 years of age; AND
- ii.** Patient has \geq 20 identifiable nodular lesions in total on both arms, and/or both legs, and/or trunk; AND
- iii.** Patient has experienced pruritus for \geq 6 weeks; AND
- iv.** Patient meets ONE of the following (a or b):
 - a)** Patient's prurigo nodularis is NOT medication-induced or secondary to a non-dermatologic condition such as neuropathy or a psychiatric disease; OR
 - b)** According to the prescriber, the patient has a secondary cause of prurigo nodularis that has been identified and adequately managed; AND
- v.** Patient meets ALL of the following (a, b, and c):

- a) Patient has tried at least one high- or super-high-potency prescription topical corticosteroid; AND
 - b) This topical corticosteroid was applied daily for at least 14 consecutive days; AND
 - c) According to the prescriber, inadequate efficacy was demonstrated with this topical corticosteroid therapy; AND
 - vi. The medication is prescribed by or in consultation with an allergist, immunologist, or dermatologist.
- B) Patient is Currently Receiving Dupixent.** Approve for 1 year if the patient meets BOTH of the following (i and ii):
- i. Patient has already received at least 6 months of therapy with Dupixent; AND
Note: A patient who has received < 6 months of therapy or who is restarting therapy with Dupixent should be considered under criterion 6A (Prurigo Nodularis, Initial Therapy).
 - ii. Patient has experienced a beneficial clinical response, defined by ONE of the following (a, b, or c):
 - (1) Reduced nodular lesion count; OR
 - (2) Decreased pruritus; OR
 - (3) Reduced nodular lesion size.

CONDITIONS NOT COVERED

- **Dupixent® (dupilumab subcutaneous injection – Regeneron/sanofi-aventis)**

is(are) considered experimental, investigational, or unproven for ANY other use(s) including the following (this list may not be all inclusive; criteria will be updated as new published data are available):

- 1. Concurrent Use of Dupixent with another Monoclonal Antibody Therapy.** The efficacy and safety of Dupixent in combination with other monoclonal antibody therapies have not been established.
Note: Monoclonal antibody therapies are Adbry® (tralokinumab-ldrm subcutaneous injection), Cinqair® (reslizumab intravenous injection), Ebglyss® (lebrikizumab-lbkz subcutaneous injection), Fasentra® (benralizumab subcutaneous injection), Nemluvio® (nemolizumab-ilto subcutaneous injection), Nucala® (mepolizumab subcutaneous injection), Tezspire® (tezepelumab-ekko subcutaneous injection), or Xolair® (omalizumab subcutaneous injection).
- 2. Concurrent Use of Dupixent with Janus Kinase (JAK) Inhibitors (oral or topical).** Use of JAK inhibitors are not recommended for use in combination with other JAK inhibitors, biologic immunomodulators (e.g., Dupixent), or with other immunosuppressants.^{29-31,35}
Note: Examples of JAK inhibitors are Cibinqo® (abrocitinib tablets), Leqselvi™ (deuruxolitinib tablets), Rinvoq®/Rinvoq® LQ (upadacitinib extended-release tablets and oral solution), and Opzelura™ (ruxolitinib cream).

REFERENCES

1. Dupixent® subcutaneous injection [prescribing information]. Tarrytown, NY: Regeneron/sanofi-aventis; September 2024.

2. Castro M, Corren J, Pavord ID, et al. Dupilumab efficacy and safety in moderate-to-severe uncontrolled asthma. *N Engl J Med*. 2018;378(26):2486-2496.
3. Wenzel S, Castro M, Corren J, et al. Dupilumab efficacy and safety in adults with uncontrolled persistent asthma despite use of medium-to-high-dose inhaled corticosteroids plus a long-acting β 2 agonist: a randomised double-blind placebo-controlled pivotal phase 2b dose-ranging trial. *Lancet*. 2016;388(10039):31-44.
4. Rabe KF, Nair P, Brusselle G, et al. Efficacy and safety of dupilumab in glucocorticoid-dependent severe asthma. *N Engl J Med*. 2018;378(26):2475-2485.
5. Bacharier LB, Maspero JF, Katelaris CH, et al. Dupilumab in children with uncontrolled moderate-to-severe asthma. *N Engl J Med*. 2021;385(24):2230-2240.
6. Simpson EL, Bieber T, Guttman-Yassky E, et al. Two phase 3 trials of dupilumab versus placebo in atopic dermatitis. *N Engl J Med*. 2016;375(24):2335-2348.
7. Blauvelt A, de Bruin-Weller M, Gooderham M, et al. Long-term management of moderate-to-severe atopic dermatitis with dupilumab and concomitant topical corticosteroids (LIBERTY AD CHRONOS): a 1-year, randomised, double-blinded, placebo-controlled, phase 3 trial. *Lancet*. 2017; 389:2287-2303.
8. Simpson EL, Paller AS, Siegfried EC, et al. Efficacy and safety of dupilumab in adolescents with uncontrolled moderate to severe atopic dermatitis: a phase 3 randomized clinical trial. *JAMA Dermatol*. 2020;156(1):44-56.
9. Paller AS, Siegfried EC, Thaci D, et al. Efficacy and safety of dupilumab with concomitant topical corticosteroids in children 6 to 11 years old with severe atopic dermatitis: a randomized, double-blinded, placebo-controlled phase 3 trial. *J Am Acad Dermatol*. 2020;83(5):1282-1293.
10. Paller AS, Simpson EL, Siegfried EC, et al. Dupilumab in children aged 6 months to younger than 6 years with uncontrolled atopic dermatitis: a randomised, double-blind, placebo-controlled, phase 3 trial. *Lancet*. 2022; 400:908-919.
11. Bachert C, Han JK, Desrosiers M, et al. Efficacy and safety of dupilumab in patients with severe chronic rhinosinusitis with nasal polyps (LIBERTY NP SINUS-24 and LIBERTY NP SINUS-52): results from two multicenter, randomized, double-blind, placebo-controlled, parallel-group phase3 trials. *Lancet*. 2019;394(10209):1638-1650.
12. Bachert C, Mannent L, Naclerio RM, et al. Effect of subcutaneous dupilumab on nasal polyp burden in patients with chronic sinusitis and nasal polyposis: a randomized clinical trial. *JAMA*. 2016;315(5):469-479.
13. Jonstam K, Swanson BN, Mannent L, et al. Dupilumab reduces local type 2 pro-inflammatory biomarkers in chronic rhinosinusitis with nasal polyposis. *Allergy*. 2019;74(4):743-752.
14. Dellon ES, Rothenberg ME, Collins MH, et al. Dupilumab in adults with adolescents with eosinophilic esophagitis. *N Engl J Med*. 2022;387(25):2317-2330.
15. Yosipovich G, Mollanazar N, Stander S, et al. Dupilumab significantly improves itch and skin lesions in patients with prurigo nodularis: results from a 2nd Phase III trial (LIBERTY-PN PRIME) [presentation]. Presented at: the European Academy of Dermatology and Venereology; Milan, Italy; September 7-10, 2022.
16. Yosipovich G, Mollanazar N, Stander S, et al. Dupilumab significantly improves itch and skin lesions in patients with prurigo nodularis: results from a Phase III trial (LIBERTY-PN PRIME 2) [presentation]. Presented at: the 2022 Annual Meeting of the American Academy of Dermatology; Boston, MA; March 25-29, 2022.
17. Global Initiative for Asthma. Global strategy for asthma management and prevention. Updated 2022. Available at: <http://www.ginasthma.org>. Accessed on April 10, 2024.
18. Chung KF, Wenzel SE, Brozek JL, et al. International ERS/ATS guidelines on definition, evaluation, and treatment of severe asthma. *Eur Respir J*. 2014; 43:343-373.
19. Holguin F, Cardet JC, Chung KF, et al. Management of severe asthma: a European Respiratory Society/American Thoracic Society Guideline. *Eur Respir J*. 2020; 55:1900588.

20. Davis DMR, Drucker AM, Alikhan A, et al. Guidelines of care for the management of atopic dermatitis in adults with phototherapy and systemic therapies. *J Am Acad Dermatol*. 2024;90(2): e43-e56.
21. Sidbury R, Alikhan A, Cohen DE, et al. Guidelines of care for the management of atopic dermatitis in adults with topical therapies. *J Am Acad Dermatol*. 2023;89(e1-e20).
22. Rank MA, Chu DK, Bognanni A, et al. Joint Task Force on Practice Parameters GRADE guidelines for the medical management of chronic rhinosinusitis with nasal polyposis. *J Allergy Clin Immunol*. 2023;151(2):386-398.
23. Peters AT, Spector S, Hsu J, et al. Diagnosis and management of rhinosinusitis: a practice parameter update. *Ann Allergy Asthma Immunol*. 2014:347-385.
24. Dykewicz MS, Wallace DV, Baroody F, et al. Treatment of seasonal allergic rhinitis: an evidenced-based focused 2017 guideline update. *Ann Allergy Asthma Immunol*. 2017;119(6):489-511.
25. Rosenfeld RM, Piccirillo JF, Chandrasekhar SS, et al. Clinical practice guideline (update): adult sinusitis. *Otolaryngol Head Neck Surg*. 2015;152(2S): S1-S39.
26. Joint Task Force on Practice Parameters: American Academy of Allergy, Asthma and Immunology and the American College of Allergy, Asthma, and Immunology. Rhinitis 2020: a practice parameter update. *J Allergy Clin Immunol*. 2020; 146:721-767.
27. Hirano I, Chan ES, Rank MA, et al. AGA Institute and Joint Task Force on Allergy-Immunology Practice Parameters Clinical Guidelines for the Management of Eosinophilic Esophagitis. *Gastroenterology*. 2020;158(6):1776-1786.
28. Elmariah S, Kim B, Berger T, et al. Practical approaches for diagnosis and management of prurigo nodularis: United States expert panel consensus. *J Am Acad Dermatol*. 2021;84(3):747-760.
29. Cibinqo® tablets [prescribing information]. New York, NY: Pfizer; February 2023.
30. Rinvoq® extended-release tablets/Rinvoq® LQ oral solution [prescribing information]. North Chicago, IL: AbbVie; April 2024.
31. Opzelura® cream [prescribing information]. Wilmington, DE: Incyte; January 2023.
32. Bhatt SP, Rabe KF, Hanania NA, et al. Dupilumab for COPD with type 2 inflammation indicated by eosinophil counts. *N Engl J Med*. 2023;289(3):205-214.
33. Bhatt SP, Rabe KF, Hanania NA, et al. Dupilumab for COPD with blood eosinophil evidence of type 2 inflammation. *N Engl J Med*. 2024;390(24):2274-2283.
34. Global Initiative for Chronic Obstructive Lung Disease. Global strategy for the diagnosis, management, and prevention of chronic obstructive pulmonary disease. National Institutes of Health, National Heart, Lung, and Blood Institute; 2024. Available at: <https://goldcopd.org/gold-reports/>. Accessed on September 30, 2024.
35. Leqselvi™ tablets [prescribing information]. Whippany, NJ: Sun/Halo; July 2024.

HISTORY

Type of Revision	Summary of Changes	Review Date
Annual Revision	Conditions Not Covered : Criteria were updated to clarify that use of Dupixent with another monoclonal antibody therapy is specific to Cinqair, Fasentra, Nucala, Tezspire, Xolair, and Adbry.	03/22/2023
Selected Revision	Conditions Not Covered : Criteria were added for "Concurrent Use of Dupixent with Janus Kinase Inhibitors (JAKis) [oral or topical]."	05/10/2023
Selected Revision	Chronic Rhinosinusitis with Nasal Polyps: Approval condition updated from "Nasal Polyps" to "Chronic	02/14/2024

	<p>Rhinosinusitis with Nasal Polyps.” Duration of the intranasal corticosteroid requirement was changed from 3 months to 4 weeks.</p> <p>Eosinophilic Esophagitis: The age of approval was reduced from ≥ 12 years of age to ≥ 1 year of age. Additionally, the weight requirement was reduced from ≥ 40 kg to ≥ 15 kg.</p>	
Annual Revision	<p>Asthma: Removed leukotriene receptor antagonists as an example of additional asthma controller or asthma maintenance medications.</p>	04/19/2024
Selected Revision	<p>Chronic Rhinosinusitis with Nasal Polyps: The age of approval was changed from ≥ 18 years of age to ≥ 12 years of age.</p> <p>Asthma: Eosinophil level requirements were clarified to require a level ≥ 150 cells/microliter either within the previous 6 weeks OR prior to treatment with a monoclonal antibody that may alter eosinophil levels. Previously, criteria required a level ≥ 150 cells/microliter either within the previous 6 weeks OR within 6 weeks prior to treatment with a monoclonal antibody that may lower eosinophil levels.</p> <p>Throughout the policy, Ebglyss (lebrikizumab-lbkz subcutaneous injection) and Nemluvio (nemolizumab-ilto subcutaneous injection) were added to notes as examples of monoclonal antibody therapies.</p>	09/25/2024
Selected Revision	<p>Chronic Obstructive Pulmonary Disease: This condition and criteria for approval were added to the policy. New approval criteria for this indication were added that include an age requirement, an eosinophil requirement, a trial of inhaled therapies, a history of chronic bronchitis signs or symptoms, a history of COPD exacerbations, and specialist involvement.</p> <p>Conditions Not Covered</p> <p>Concurrent Use of Dupixent with Janus Kinase (JAK) Inhibitors (oral or topical): Leqselvi™ (deuruxolitinib tablets) and Rinvoq® LQ (upadacitinib oral solution) were added as examples of JAK inhibitors.</p>	10/09/2024

"Cigna Companies" refers to operating subsidiaries of The Cigna Group. All products and services are provided exclusively by or through such operating subsidiaries, including Cigna Health and Life Insurance Company, Connecticut General Life Insurance Company, Evernorth Behavioral Health, Inc., Cigna Health Management, Inc., and HMO or service company subsidiaries of The Cigna Group. © 2024 The Cigna Group.