



PRIOR AUTHORIZATION POLICY

POLICY: Oncology – Imatinib Prior Authorization Policy

- Gleevec® (imatinib tablets– Novartis, generic)

REVIEW DATE: 05/01/2024

INSTRUCTIONS FOR USE

THE FOLLOWING COVERAGE POLICY APPLIES TO HEALTH BENEFIT PLANS ADMINISTERED BY CIGNA COMPANIES. CERTAIN CIGNA COMPANIES AND/OR LINES OF BUSINESS ONLY PROVIDE UTILIZATION REVIEW SERVICES TO CLIENTS AND DO NOT MAKE COVERAGE DETERMINATIONS. REFERENCES TO STANDARD BENEFIT PLAN LANGUAGE AND COVERAGE DETERMINATIONS DO NOT APPLY TO THOSE CLIENTS. COVERAGE POLICIES ARE INTENDED TO PROVIDE GUIDANCE IN INTERPRETING CERTAIN STANDARD BENEFIT PLANS ADMINISTERED BY CIGNA COMPANIES. PLEASE NOTE, THE TERMS OF A CUSTOMER'S PARTICULAR BENEFIT PLAN DOCUMENT [GROUP SERVICE AGREEMENT, EVIDENCE OF COVERAGE, CERTIFICATE OF COVERAGE, SUMMARY PLAN DESCRIPTION (SPD) OR SIMILAR PLAN DOCUMENT] MAY DIFFER SIGNIFICANTLY FROM THE STANDARD BENEFIT PLANS UPON WHICH THESE COVERAGE POLICIES ARE BASED. FOR EXAMPLE, A CUSTOMER'S BENEFIT PLAN DOCUMENT MAY CONTAIN A SPECIFIC EXCLUSION RELATED TO A TOPIC ADDRESSED IN A COVERAGE POLICY. IN THE EVENT OF A CONFLICT, A CUSTOMER'S BENEFIT PLAN DOCUMENT ALWAYS SUPERSEDES THE INFORMATION IN THE COVERAGE POLICIES. IN THE ABSENCE OF A CONTROLLING FEDERAL OR STATE COVERAGE MANDATE, BENEFITS ARE ULTIMATELY DETERMINED BY THE TERMS OF THE APPLICABLE BENEFIT PLAN DOCUMENT. COVERAGE DETERMINATIONS IN EACH SPECIFIC INSTANCE REQUIRE CONSIDERATION OF 1) THE TERMS OF THE APPLICABLE BENEFIT PLAN DOCUMENT IN EFFECT ON THE DATE OF SERVICE; 2) ANY APPLICABLE LAWS/REGULATIONS; 3) ANY RELEVANT COLLATERAL SOURCE MATERIALS INCLUDING COVERAGE POLICIES AND; 4) THE SPECIFIC FACTS OF THE PARTICULAR SITUATION. EACH COVERAGE REQUEST SHOULD BE REVIEWED ON ITS OWN MERITS. MEDICAL DIRECTORS ARE EXPECTED TO EXERCISE CLINICAL JUDGMENT AND HAVE DISCRETION IN MAKING INDIVIDUAL COVERAGE DETERMINATIONS. COVERAGE POLICIES RELATE EXCLUSIVELY TO THE ADMINISTRATION OF HEALTH BENEFIT PLANS. COVERAGE POLICIES ARE NOT RECOMMENDATIONS FOR TREATMENT AND SHOULD NEVER BE USED AS TREATMENT GUIDELINES. IN CERTAIN MARKETS, DELEGATED VENDOR GUIDELINES MAY BE USED TO SUPPORT MEDICAL NECESSITY AND OTHER COVERAGE DETERMINATIONS.

CIGNA NATIONAL FORMULARY COVERAGE:

OVERVIEW

Imatinib, a tyrosine kinase inhibitor (TKI), is indicated for the following uses:^{1,2}

- **Acute lymphoblastic leukemia (ALL)**, Philadelphia chromosome positive (Ph+), in adults with relapsed or refractory disease.
- **ALL**, newly diagnosed and Ph+, in combination with chemotherapy in pediatric patients.
- **Aggressive systemic mastocytosis**, without the D816V c-Kit mutation or with unknown c-Kit mutational status, in adults.
- **Chronic myeloid leukemia (CML)**, newly diagnosed and Ph+, chronic phase in adult and pediatric patients.
- **CML**, Ph+, in blast phase, accelerated phase, or in chronic phase in patients after failure of interferon alfa therapy.
- **Dermatofibrosarcoma protuberans** in adults with unresectable, current, and/or metastatic disease.
- **Gastrointestinal stromal tumors (GIST)**, in patients with KIT (CD117) positive unresectable and/or metastatic malignant disease.
- **GIST**, Kit (CD117) positive, as adjuvant treatment of adults following resection.
- **Hypereosinophilic syndrome and/or chronic eosinophilic leukemia**, in adults who have the *FIP1L1-PDGFR* alpha fusion kinase (mutation analysis or

fluorescence in situ hybridization demonstration of CICH2 allele deletion) and for patients with hypereosinophilic syndrome and/or chronic eosinophilic leukemia who are *FIP111-PDGFR* alpha fusion kinase negative or unknown.

- **Myelodysplastic/myeloproliferative diseases**, associated with platelet-derived growth factor receptor (*PDGFR*) gene rearrangements in adults.

Guidelines

Imatinib is addressed in guidelines from National Comprehensive Cancer Network (NCCN):

- **ALL:** NCCN guidelines for adults and adolescents (version 4.2023 – February 5, 2024) recommend imatinib for Ph+ disease in many different clinical circumstances (e.g., induction, consolidation therapy, maintenance, or relapsed or refractory disease) [category 2A].³ NCCN guidelines for pediatric ALL (version 5.2024 – April 3, 2024) feature imatinib prominently (category 2A) in a variety of clinical scenarios.⁴
- **Bone Cancer:** NCCN guidelines (version 2.2024 – March 12, 2024) recommend imatinib either as monotherapy or as “other recommended regimens” or in combination with cisplatin or Rapamune® (sirolimus tablets) for chordoma as “useful in certain circumstances” (both category 2A).⁵
- **CML:** NCCN guidelines (version 2.2024 – December 5, 2023) state that for patients with chronic phase CML with a low-risk score, the primary “preferred” treatment recommendations includes a first-generation TKI (imatinib) or a second-generation TKI (Bosulif® [bosutinib tablets], Sprycel® [dasatinib tablets], or Tassigna® [nilotinib capsules] {all category 1}).⁶ For patients with chronic phase CML with an intermediate- or high-risk score, a second-generation TKI is preferred (Bosulif, Sprycel, or Tassigna [all category 1]); imatinib is listed as “other recommended regimen” (category 2A); imatinib is also recommended for other clinical scenarios (category 2A).
- **Dermatofibrosarcoma Protuberans:** NCCN guidelines (version 1.2024 – November 9, 2023) state to consider neoadjuvant imatinib for unresectable disease, borderline resectable disease, and in patients whom resection with negative margins may result in unacceptable functional or cosmetic outcomes (category 2A) and for recurrent disease in cases where the disease is unresectable, or unacceptable functional or adverse cosmetic outcomes may occur with resection (category 2A).⁷
- **GIST:** NCCN guidelines (version 1.2024 – March 8, 2024) recommend imatinib as a “preferred regimen” for first-line therapy (category 1) in various scenarios (e.g., for sensitive mutations or for *PDGFRA* exon 18 mutations [excluding the D842V mutation) and is recommended in other clinical scenarios (e.g., neoadjuvant and adjuvant therapy)[category 2A].⁸
- **Graft-Versus-Host Disease (GVHD):** NCCN guidelines for hematopoietic cell transplantation (version 1.2024 – April 26, 2024) address GVHD.⁹ Imatinib is cited as one of many therapies recommended for steroid-refractory, chronic GVHD (category 2A).
- **Kaposi Sarcoma:** NCCN guidelines (version 1.2024 – November 7, 2023) recommend imatinib for subsequent systemic therapy for relapsed/refractory therapy as “useful in certain circumstances.”¹⁰ First-line systemic therapy options are liposomal doxorubicin as “preferred regimen” and paclitaxel.

- **Melanoma: Cutaneous:** NCCN guidelines (version 2.2024 – April 3, 2024) recommend imatinib as second-line or subsequent therapy for metastatic or unresectable disease for tumors with activating mutations of *KIT* as “useful in certain circumstances” (category 2A).¹¹
- **Myelodysplastic Syndromes:** NCCN guidelines (version 1.2024 – February 12, 2024) note that data have demonstrated that patients with chronic myelomonocytic leukemia who have *PDGFRβ* may respond well to imatinib.¹²
- **Myeloid/Lymphoid Neoplasms with Eosinophilia and Tyrosine Kinase Gene Fusions:** NCCN guidelines (version 1.2024 – December 21, 2023) recommend imatinib for patients with *ABL1* rearrangements in the chronic phase or blast phase (category 2A).¹³ Imatinib is also recommended for certain situations where the tumor has an *FIP1L1-PDGFRα* or *PDGFRβ* rearrangement (category 2A). Imatinib is also recommended for treatment in combination with ALL- or acute myeloid leukemia-type induction chemotherapy followed by allogeneic hematopoietic stem cell transplantation (if eligible) for lymphoid, myeloid, or mixed lineage neoplasms with eosinophilia and *ABL1* rearrangement in blast phase (category 2A).
- **Soft Tissue Sarcomas:** NCCN guidelines (version 1.2024 – April 26, 2024) recommend imatinib for desmoid tumors (aggressive fibromatosis) as a “preferred regimen” (category 2A). For dermatofibrosarcoma protuberans with fibrosarcomatous transformation, imatinib is recommended as a “preferred regimen” (category 2A). For pigmented villonodular synovitis/tenosynovial giant cell tumor, imatinib is recommended as “useful in certain circumstances” (category 2A).¹⁴
- **Systemic Mastocytosis:** NCCN guidelines (version 1.2023 – May 24, 2023,) recommend imatinib for aggressive systemic mastocytosis (for *KIT* D816V mutation negative or unknown; well differentiated systemic mastocytosis; eosinophilia is present with *FIP1L1-PDGFRα* fusion gene) as “useful in certain circumstances” (category 2A).¹⁵

POLICY STATEMENT

Prior Authorization is recommended for prescription benefit coverage of imatinib tablets. All approvals are provided for the duration noted below.

- **Gleevec® (imatinib tablets(Novartis, generic)**

is(are) covered as medically necessary when the following criteria is(are) met for FDA-approved indication(s) or other uses with supportive evidence (if applicable):

FDA-Approved Indications

- 1. Acute Lymphoblastic Leukemia.** Approve for 1 year if the patient has Philadelphia chromosome-positive acute lymphoblastic leukemia.
- 2. Aggressive Systemic Mastocytosis.** Approve for 1 year if the patient is ≥ 18 years of age.

- 3. Chronic Myeloid Leukemia.** Approve for 1 year if the patient has Philadelphia chromosome-positive chronic myeloid leukemia.
- 4. Dermatofibrosarcoma Protuberans.** Approve for 1 year if the patient is ≥ 18 years of age.
- 5. Gastrointestinal Stromal Tumors.** Approve for 1 year.
- 6. Hypereosinophilic Syndrome and/or Chronic Eosinophilic Leukemia.** Approve for 1 year if the patient is ≥ 18 years of age.
- 7. Myelodysplastic/Myeloproliferative Disease.** Approve for 1 year if the patient meets BOTH of the following (A and B):
 - A) Patient is ≥ 18 years of age; AND
 - B) The condition is associated with platelet-derived growth factor receptor (*PDGFR*) gene rearrangements.

Other Uses with Supportive Evidence

- 8. Chordoma.** Approve for 1 year.
- 9. Desmoid Tumors (Aggressive Fibromatosis).** Approve for 1 year
- 10. Graft-Versus-Host Disease, Chronic.** Approve for 1 year if the patient has tried at least one conventional systemic treatment for graft-versus-host disease.
Note: Examples include corticosteroids (methylprednisolone, prednisone); cyclosporine; tacrolimus; mycophenolate mofetil; Imbruvica (ibrutinib capsules, tablets, and oral suspension); low-dose methotrexate; sirolimus; Rezurock (belumosudil tablets); and Jakafi (ruxolitinib tablets).
- 11. Kaposi Sarcoma.** Approve for 1 year if the patient meets ALL of the following (A, B and C):
 - A)** Patient is ≥ 18 years of age; AND
 - B)** Patient has tried at least one medication; AND
Note: Examples include liposomal doxorubicin, paclitaxel, Pomalyst (pomalidomide capsules), lenalidomide, etoposide, and Thalomid (thalidomide capsules).
 - C)** Patient has relapsed or refractory disease.
- 12. Melanoma, Cutaneous.** Approve for 1 year if the patient meets ALL of the following (A, B, C, and D):
 - A)** Patient is ≥ 18 years of age; AND
 - B)** Patient has metastatic or unresectable disease; AND
 - C)** Patient has an activating *KIT* mutation; AND
 - D)** Patient has tried at least one systemic regimen.
Note: Examples of a systemic regimen include: Opdivo (nivolumab intravenous infusion) + Yervoy (ipilimumab intravenous infusion), Opdivo + Opdualag (nivolumab/relatlimab-rmbw intravenous infusion), Keytruda

(pembrolizumab intravenous infusion), Opdivo, Tafinlar (dabrafenib capsules and tablets for suspension) + Mekinist (trametinib tablets), Zelboraf (vemurafenib tablets) + Cotellic (cobimetinib tablets), Braftovi (encorafenib capsules) + Mektovi (binimetinib tablets).

13. Myeloid/Lymphoid Neoplasms with Eosinophilia. Approve for 1 year if the patient meets BOTH of the following (A and B):

A) Patient is \geq 18 years of age; AND

B) Patient meets ONE of the following (i or ii):

i. The tumor has an *ABL1* rearrangement; OR

ii. The tumor has an *FIP1L1-PDGFR* or *PDGFRB* rearrangement.

14. Pigmented Villonodular Synovitis/Tenosynovial Giant Cell Tumor.

Approve for 1 year if the patient meets ONE of the following (A or B):

A) Patient has tried Turalio (pexidartinib capsules); OR

B) Patient cannot take Turalio, according to the prescriber.

Note: Examples of reasons for not being able to take Turalio include patients with elevated liver enzymes or concomitant use of medications that are associated with hepatotoxicity.

CONDITIONS NOT COVERED

- **Gleevec® (imatinib tablets(Novartis, generic)**

is(are) considered experimental, investigational, or unproven for ANY other use(s).

REFERENCES

1. Gleevec® tablets [prescribing information]. East Hanover, NJ: Novartis; March 2024.
2. Imatinib tablets [prescribing information]. Cranbury, NJ: Sun; Sept 2022.
3. The NCCN Acute Lymphoblastic Leukemia Clinical Practice Guidelines in Oncology (version 4.2023 – February 5, 2024). © 2024 National Comprehensive Cancer Network. Available at: <http://www.nccn.org>. Accessed on April 29, 2024.
4. The NCCN Pediatric Acute Lymphoblastic Leukemia Clinical Practice Guidelines in Oncology (version 5.2024 – April 3, 2024). © 2024 National Comprehensive Cancer Network. Available at: <http://www.nccn.org>. Accessed on April 29, 2024.
5. The NCCN Bone Cancer Clinical Practice Guidelines in Oncology (version 2.2024 – March 12, 2024). © 2024 National Comprehensive Cancer Network. Available at: <http://www.nccn.org>. Accessed on April 29, 2024.
6. The NCCN Chronic Myeloid Leukemia Clinical Practice Guidelines in Oncology (version 2.2024 – December 5, 2023). © 2023 National Comprehensive Cancer Network. Available at: <http://www.nccn.org>. Accessed on April 29, 2024.
7. The NCCN Dermatofibrosarcoma Protuberans Clinical Practice Guidelines in Oncology (version 1.2024 – November 9, 2023) © 2023 National Comprehensive Cancer Network. Available at: <http://www.nccn.org>. Accessed on April 29, 2024.
8. The NCCN Gastrointestinal Stromal Tumors Clinical Practice Guidelines in Oncology (version 1.2024 – March 8, 2024). © 2024 National Comprehensive Cancer Network. Available at: <http://www.nccn.org>. Accessed on April 29, 2024.
9. The NCCN Hematopoietic Cell Transplantation (HCT) Clinical Practice Guidelines in Oncology (version 1.20234 – April. 26, 2024). © 2024 National Comprehensive Cancer Network. Available at: <http://www.nccn.org>. Accessed on April 29, 2024.

10. The NCCN Kaposi Sarcoma Clinical Practice Guidelines in Oncology (version 1.2024 – November 7, 2023). © 2023 National Comprehensive Cancer Network. Available at: <http://www.nccn.org>. Accessed on April 29, 2024.
11. The NCCN Melanoma: Cutaneous Clinical Practice Guidelines in Oncology (version 2.2024 – April 3, 2024). © 2024 National Comprehensive Cancer Network. Available at: <http://www.nccn.org>. Accessed on April 29, 2024.
12. The NCCN Myelodysplastic Syndromes Clinical Practice Guidelines in Oncology (version 1.2024 – February 12, 2024). © 2024 National Comprehensive Cancer Network. Available at: <http://www.nccn.org>. Accessed on April 29, 2024.
13. The NCCN Myeloid/Lymphoid Neoplasms with Eosinophilia and Tyrosine Kinase Gene Fusions Clinical Practice Guidelines in Oncology (version 1.2024 – December 21, 2023). © 2023 National Comprehensive Cancer Network. Available at: <http://www.nccn.org>. Accessed on April 29, 2024.
14. The NCCN Soft Tissue Sarcoma Clinical Practice Guidelines in Oncology (version 1.2024 – April 26, 2024). © 2024 National Comprehensive Cancer Network. Available at: <http://www.nccn.org>. Accessed on April 29, 2024.
15. The NCCN Systemic Mastocytosis Clinical Practice Guidelines in Oncology (version 3.2024 – April 24, 2024). © 2024 National Comprehensive Cancer Network. Available at: <http://www.nccn.org>. Accessed on April 29, 2024.

HISTORY

Type of Revision	Summary of Changes	Review Date
Annual Revision	<p>Desmoid tumors (aggressive fibromatosis): The “fibromatosis (desmoid tumors) indication was reworded to “Desmoid tumors (aggressive fibromatosis).” The requirement that the patient has advanced or unresectable disease was removed.</p> <p>Kaposi Sarcoma: The requirement that the patient is ≥ 18 years of age was added.</p> <p>Melanoma, Cutaneous: The “metastatic melanoma” indication was reworded to “melanoma, cutaneous.” The criterion that the patient has “advanced/recurrent” disease was changed to “unresectable.” The criterion that the patient has “c-kit-positive melanoma” was reworded to patient has an “activating <i>KIT</i> mutation.” The following criteria were added: Patient is ≥ 18 years of age; and patient has tried at least one systemic regimen with a note with examples of a systemic regimen.</p>	05/31/2023
Annual Revision	No criteria change.	05/01/2024

"Cigna Companies" refers to operating subsidiaries of The Cigna Group. All products and services are provided exclusively by or through such operating subsidiaries, including Cigna Health and Life Insurance Company, Connecticut General Life Insurance Company, Evernorth Behavioral Health, Inc., Cigna Health Management, Inc., and HMO or service company subsidiaries of The Cigna Group. © 2024 The Cigna Group.