

RETIRED

Valid for dates of service prior to 11/1/24 only
For dates of service 11/1 and after, see policy:

[EviCore Cigna Commercial Membership | EviCore by Evernorth](#)



Medical Coverage Policy

Effective Date 11/15/2023
Next Review Date 11/15/2024
Coverage Policy Number 0393

Discography

Table of Contents

Overview.....	2
Coverage Policy.....	2
General Background.....	3
Medicare Coverage Determinations.....	10
Coding Information.....	10
References.....	11
Revision Details.....	17

Related Coverage Resources

[Intervertebral Disc \(IVD\) Prostheses](#)
[Lumbar Fusion for Spinal Instability and Degenerative Disc Conditions, Including Sacroiliac Fusion](#)
[Minimally Invasive Spine Surgery Procedures and Trigger Point Injections](#)
[Spine Imaging Guidelines](#)

RETIRED

Valid for dates of service prior to 11/1/24 only

For dates of service 11/1 and after, see policy:

[EviCore Cigna Commercial Membership | EviCore by Evernorth](#)

INSTRUCTIONS FOR USE

The following Coverage Policy applies to health benefit plans administered by Cigna Companies. Certain Cigna Companies and/or lines of business only provide utilization review services to clients and do not make coverage determinations. References to standard benefit plan language and coverage determinations do not apply to those clients. Coverage Policies are intended to provide guidance in interpreting certain standard benefit plans administered by Cigna Companies. Please note, the terms of a customer's particular benefit plan document [Group Service Agreement, Evidence of Coverage, Certificate of Coverage, Summary Plan Description (SPD) or similar plan document] may differ significantly from the standard benefit plans upon which these Coverage Policies are based. For example, a customer's benefit plan document may contain a specific exclusion related to a topic addressed in a Coverage Policy. In the event of a conflict, a customer's benefit plan document always supersedes the information in the Coverage Policies. In the absence of a controlling federal or state coverage mandate, benefits are ultimately determined by the terms of the applicable benefit plan document. Coverage determinations in each specific instance require consideration of 1) the terms of the applicable benefit plan document in effect on the date of service; 2) any applicable laws/regulations; 3) any relevant collateral source materials including Coverage Policies and; 4) the specific facts of the particular situation. Each coverage request should be reviewed on its own merits. Medical directors are expected to exercise clinical judgment where appropriate and have discretion in making individual coverage determinations. Where coverage for care or services does not depend on specific circumstances, reimbursement will only be provided if a requested service(s) is submitted in accordance with the relevant criteria outlined in the applicable Coverage Policy, including covered diagnosis and/or procedure code(s). Reimbursement is not allowed for services when billed for conditions or diagnoses that are not covered under this Coverage Policy (see "Coding Information" below). When billing, providers must use the most appropriate codes as of the effective date of the submission. Claims submitted for services that are not accompanied by covered code(s) under the applicable Coverage Policy will be denied as not covered. Coverage Policies relate exclusively to the administration of health benefit plans. Coverage Policies are not recommendations for treatment and should never be used as treatment guidelines. In certain markets, delegated vendor guidelines may be used to support medical necessity and other coverage determinations.

Overview

This Coverage Policy addresses discography, also referred to as stimulation discography, or a discogram, an imaging test used to determine if a specific disc is causing back pain.

Coverage Policy

Lumbar provocative discography, including post discography computed tomography (CT) assessment, is considered medically necessary for the preoperative evaluation of discogenic back pain, when ALL of the following criteria for a single level lumbar fusion have been met:

- unremitting pain with significant functional impairment of at least twelve months duration
- failure of at least six (6) consecutive months of structured*, physician-supervised conservative medical management, which includes **ALL** of the following components:
 - exercise, including core stabilization exercises
 - nonsteroidal and/or steroidal medication (unless contraindicated)
 - physical therapy, including passive and active treatment modalities
 - activity/lifestyle modification
 - participation in three (3) or more individual or group cognitive behavioral

RETIRED

Valid for dates of service prior to 11/1/24 only

For dates of service 11/1 and after, see policy:

[EviCore Cigna Commercial Membership | EviCore by Evernorth](#)

therapy (CBT) sessions provided by a licensed healthcare professional, with competence in principles and practice of CBT, (e.g., PT, OT, psychiatrist, psychologist, social worker, psychiatric nurse, other licensed professional) that include ALL of the following elements:

- disease education
- activity and lifestyle modification
- stress management (stress management typically also includes strategies to deal with emotions such as fear, anxiety, sadness that can interfere with pain management)
- complex imaging studies (e.g., computed tomography [CT] scan or magnetic resonance imaging [MRI] scan) do not conclusively demonstrate single level degenerative disc disease as the likely cause of pain
- statement from a primary care physician, neurologist, physiatrist, psychiatrist, psychologist, or other licensed behavioral and/or other medical healthcare provider attesting to the absence of untreated, underlying mental health conditions/issues (e.g., depression, drug and alcohol abuse) as a major contributor to chronic pain

***Note:** Structured medical management consists of medical care that is delivered through regularly scheduled appointments, including follow-up evaluation, with licensed healthcare professionals.

Lumbar discography (e.g., provocative discography, stimulation discography) with or without CT assessment, is considered not medically necessary when performed in connection with ANY other procedure in the lumbar spine, OR in anticipation of any procedure that has been determined to be experimental, investigational or unproven, including, but not limited to ANY of the following:

RETIRED

Valid for dates of service prior to 11/1/24 only

For dates of service 11/1 and after, see policy:

[EviCore Cigna Commercial Membership | EviCore by Evernorth](#)

- intradiscal electrothermal therapy (IDET™)
- disc nucleoplasty, decompression nucleoplasty, or Coblation® Nucleoplasty™
- laser discectomy (e.g., percutaneous, laparoscopic)
- intradiscal injections (e.g., platelet rich plasma, steroid, hyaluronic acid)

Lumbar provocative discography, with or without CT assessment, is considered not medically necessary for ANY other indication.

The following discography procedures are each considered experimental, investigational or unproven:

- cervical discography
- thoracic discography
- functional anesthetic discography (FAD)
- contrast disc analysis mapping

General Background

Low back pain is the most common musculoskeletal problem in the world, and the leading cause of activity limitation and work absenteeism. It is estimated that globally, over 577 million people experience low back pain, and that it is more prevalent in women compared to men (8.01% versus 7.47%, respectively). Low back pain is found in individuals of all ages, and peaks between the ages of 45 and 49 (Wu, et al., 2020a; Duthey, 2013). Risk factors for chronic back pain include obesity, smoking, and a sedentary lifestyle. Persons with lower socioeconomic, employment, and/or educational status are also at a higher risk for chronic low back pain (Karran, et al., 2020).

Provocative Discography

Provocative discography (or discogram) is a minimally invasive diagnostic procedure used to help determine if a person's back pain is caused by a specific disc (the cushioning pads that separate the vertebrae of the spine). It is typically performed on an individual with chronic axial (i.e., mechanical, localized) back pain after other diagnostic interventions have failed to definitively identify the source of the pain, and when surgery is being considered. The procedure involves inserting a needle into the nucleus pulposus of the suspected painful disc under fluoroscopic guidance, followed by the injection of contrast dye into the disc. The injection of contrast dye allows for the determination of pressures within the disc, to help determine if the disc is the cause of the pain. Additional information regarding the shape and integrity of the disc can be obtained by fluoroscopy at the time of injection or with post-injection computed tomography (CT).

It is believed that discography induces pain as a result of the increased pressure from the contrast dye, neurochemical stimulation, and/or an increase in intervertebral pressure. A discogram is considered positive (i.e., an abnormal discogram) if, when the dye is injected into the disc, the individual reports pain that is similar to or exactly like their chronic back pain, accompanied by abnormal imaging of the disc (Gardocki and Park, 2021). These imaging patterns may be described as cotton ball, lobular, irregular, fissured, or ruptured. Cotton ball and lobular imaging patterns are typically considered normal, while irregular, fissured, and ruptured patterns indicate progressive disc degeneration (Devlin, 2021). Some published studies differ in the definition of an abnormal discogram; as a result, the technique and interpretation of the discogram must also be closely evaluated when assessing outcomes. If the discogram does not cause pain or produces pain that is not comparable with the individual's usual pain, then the cause of the pain cannot be attributed to that particular disc.

RETIRED

Valid for dates of service prior to 11/1/24 only

For dates of service 11/1 and after, see policy:

[EviCore Cigna Commercial Membership | EviCore by Evernorth](#)

Discography is often performed to assess whether an individual with painful degenerative disc disease would benefit from single level lumbar fusion. As such, consideration is given to whether the individual is an appropriate candidate for lumbar fusion surgery. Multiple factors are taken into account, including length and severity of back pain, resulting functional impairment, and persistence of symptoms despite a trial of supervised conservative medical management. Conservative management may include exercise, physical therapy, steroidal/nonsteroidal medications, lifestyle modification, and cognitive behavioral therapy (NASS, 2021; Cherkin and Herman, 2018; Brox, et al., 2010; Fairbank, et al., 2005). Please reference Cigna Medical Coverage policy "Lumbar Fusion for Spinal Instability and Degenerative Disc Conditions, Including Sacroiliac Fusion" for additional information.

Discography is typically not used as a baseline imaging study. In clinical practice, a majority of discograms are performed to evaluate the lower three lumbar discs. Lumbar discography is particularly useful as a test to exclude levels from operative intervention. Surgery may be precluded by failing to find a painful disc on discography, finding multiple painful discs, or indeterminate results (Thiyagarajah, et al., 2021). Provocative discography is less frequently performed to assess cervical or thoracic disc pain. The approach for thoracic discography is similar to the lumbar, while cervical discography is approached anteriorly rather than posteriorly.

Discography has been proven to be a safe procedure although there are associated risks. Risks and complications include disc space infection (discitis), nerve root injury, subarachnoid puncture, urticaria, bleeding, nucleus pulposus pulmonary embolism, nausea, increased pain, and spinal headache. Acute lumbar disc herniation as a result of discography has been reported (Poynton, et al., 2005). Thoracic discography also has the inherent risks of pneumothorax and direct spinal cord trauma. Discography is contraindicated for patients with dye sensitivity, spinal cord compression, prior fusion at the level being studied, or who have current local infections (Gardocki and Park, 2021; Stretanski and Vu, 2023). While older studies reported findings of disc disease progression post-discography, results from more recent studies have shown discography does not lead to acceleration of degenerative disc disease or disc herniation (Karaarslan, et al., 2019; McCormick, et al., 2019).

Discography has been proposed for use prior to or in conjunction with various minimally invasive procedures to treat back pain, including intradiscal electrothermal therapy (IDET), nucleoplasty, laser discectomy, and intradiscal injections. Presently there is insufficient evidence in the published medical literature to demonstrate the safety, efficacy, and long term outcomes of these procedures. Please reference Cigna Medical Coverage policy "Minimally Invasive Spine Surgery Procedures and Trigger Point Injections" for additional information.

U.S. Food and Drug Administration (FDA): Discography is a provocative imaging study and thus the procedure itself is not regulated by the FDA. However, several needles and catheters used to access the intervertebral disc during a discogram have been cleared via the FDA 510(k) premarket notification process, as Class II devices. Examples include the DiscCath Needle Set (DiscCath LLC, New York, NY; Sept 2021), and the Pakter Curved Needle Set (Cook Incorporated, Bloomington, IN; Jun 2018).

Cervical Discography

Cervical discography requires an anterior approach and has been recommended for patients with persistent neck pain without localized neurological findings when standard imaging studies are negative. Potential complications are related to the surrounding anatomy and include injury to the trachea, esophagus, carotid artery, discitis, spinal cord injury and pneumothorax. There are no morphologic features apparent on discography of the cervical spine that help make a diagnosis of discogenic pain. The diagnosis relies entirely on the patient's report of concordant pain, and non-

RETIRED

Valid for dates of service prior to 11/1/24 only

For dates of service 11/1 and after, see policy:

[EviCore Cigna Commercial Membership | EviCore by Evernorth](#)

painful control discs must be present. There is a paucity of literature addressing the correlation of imaging findings with cervical discography (Diehn, et al., 2023).

Literature Review: There is limited published evidence evaluating cervical discography. While there is some evidence in the form of case series and systematic reviews to support some utility for cervical discography, evidence from well-designed controlled trials is lacking. There is much debate regarding false-positive results and concerns regarding safety. The current evidence in the published peer-reviewed scientific literature is insufficient and does not lead to strong conclusions regarding clinical utility. Furthermore, what effect, if any, cervical discography has on surgical treatment for discogenic type pain has yet to be proven (Onyewu, et al., 2012; Buenaventura, et al., 2007; Shah, et al., 2005; Zheng, et al., 2004; Grubb and Kelly, 2000; Ohnmeiss, et al., 2000).

Manchikanti et al. (2018) published a systematic review of the diagnostic accuracy of lumbar, cervical, and thoracic discography. As part of the review the authors reviewed eight studies which met their inclusion criteria, five assessed lumbar discography (strength of evidence level 3) and three studies assessed cervical discography (strength of evidence level 4). For cervical and thoracic discography the available literature and value and validity continued to be low.

The results of a systematic review evaluating cervical discography concluded that despite a paucity of evidence and discrepancy among studies, the diagnostic accuracy of cervical discography has moderate validity and moderate predictive value based on modified United States Preventive Services Task Force (USPSTF) criteria (Manchikanti, et al., 2009a). This conclusion was based on Level II-2 evidence which is evidence obtained from at least one properly designed small diagnostic accuracy study. For this systematic review, the authors reviewed a total of 33 studies, three studies which met the inclusion criteria utilizing IASP (International Association for the Study of Pain) criteria: provocation discography with control discs and involving patients with chronic pain of at least three months duration.

Thoracic Discography

Thoracic discography is considered by some providers to be useful in clinical practice for the assessment of thoracic, chest, and upper abdominal pain. Similar to cervical discography, potential complications are related to the surrounding anatomy of the thoracic spine and include pneumothorax, spinal trauma, discitis and bleeding.

Literature Review: The evidence supporting the safety and utility of thoracic discography is limited, consisting mainly of case series and systematic reviews (Buenaventura, et al., 2007; Shah, et al., 2005; Wood, et al., 1999). Within a systematic review published by Manchikanti et al. (2018) evaluating lumbar, cervical and thoracic discography, the authors reported the evidence for thoracic discography was nonexistent. Included in the authors search were only studies utilizing controlled discography with IASP standards or analgesic discography, and only the studies with appropriate assessment and statistical evaluation for diagnostic prevalence. Similar to cervical discography, the limited evidence evaluating thoracic discography does not lead to strong conclusions regarding safety and clinical utility. Currently the evidence is insufficient to support the clinical value of thoracic discography.

Lumbar Discography

The clinical value of lumbar discography has been widely debated. The diagnosis of discogenic pain due to disc degeneration, internal disc disruption or annular tears, for example, is considered difficult by many authors. Not all patients with disc disease experience pain. In addition, it is reported that the ability of patients to separate spinal pain from nonspinal sources of lumbar pain may be questionable (Carragee, et al., 1999). Early published data showed high (37%) false-positive rates in a group of asymptomatic patients; however, more recent authors have reported

RETIRED

Valid for dates of service prior to 11/1/24 only

For dates of service 11/1 and after, see policy:

[EviCore Cigna Commercial Membership | EviCore by Evernorth](#)

lower false-positive rates. Several systematic reviews and case studies have been published evaluating lumbar discography. Although there is no general consensus regarding validity of discography as a diagnostic tool, authors generally agree that lumbar discography is an appropriate diagnostic test with some clinical utility for patients with low back pain, particularly when lumbar surgery is being considered, and when noninvasive diagnostic tests are inconclusive. Further, advances in procedural technique, standardized protocol, and patient selection have significantly reduced procedural risks and rates of false positive results (Gardocki and Park, 2021).

The medical literature suggests the predictive value of provocative discography on surgical outcomes has not yet been firmly established. Discography is often performed prior to arthrodesis, minimally invasive surgery and other intradiscal surgeries. Arthrodesis (spinal fusion) is a surgical method of controlling low back pain attributable to abnormal or unstable vertebrae and pain due to mechanical degeneration and is indicated when degenerative disc disease is limited to a single level. However it has been reported there was no difference in operative outcome between groups of subjects who had preoperative discography and those who did not (Madan, et al., 2002). Discography has also been recommended for use prior to minimally invasive surgeries for patients with contained disc herniation; discography may define disc containment for those candidates (Guyer and Ohnmeiss, 2003). Discography is also often performed in combination with intradiscal electrothermal therapy (IDET) as a treatment for disc pain; however, the long-term results of IDET remain unknown.

Literature Review: Evidence in the published medical literature evaluating lumbar discography consists of systemic reviews, meta-analyses, technology assessments, prospective and retrospective clinical trials, randomized trials, and published reviews (Wu, et al., 2020b; Xi, et al., 2016; Margetic, et al., 2013; Manchikanti, et al., 2013a; Manchikanti, et al., 2013b; Manchikanti, et al., 2009a; Wolfer, et al., 2008; Buenaventura, et al., 2007; Carragee, et al., 2006; Shah, et al., 2005; Cohen, et al., 2005; Guyer and Ohnmeiss, 2003). Many studies lack control groups and randomization. There is no general consensus regarding the clinical utility of lumbar discography; some authors have evaluated diagnostic accuracy and reported the evidence is strong to moderate for discography as an imaging tool for the evaluation of chronic low back pain (Manchikanti, et al., 2009a; Buenaventura, et al., 2007; Shah, et al., 2005) although Carragee et al. (2006) reported diagnostic accuracy is not high and discography has a calculated positive predictive value of 50-60%. Cohen et al. (2005) reported discography is less accurate than MRI in diagnosing herniated nucleus pulposus, although comparable or slightly more sensitive in detecting degenerative disc disease. As demonstrated in a systematic review by Manchikanti et al., (2013b) the evidence supporting diagnostic accuracy of provocation discography, after controlling for various factors which included methodological flaws, lack of standardization, and the absence of well-designed studies, is fair (according to USPSTF criteria: [good, fair, limited or poor]). According to a systematic review published by Manchikanti et al. (2018) "lumbar provocation discography performed according to the IASP criteria may be a useful tool for evaluating chronic lumbar discogenic pain. Based on modified best evidence synthesis, the indicated strength of evidence was Level III for lumbar discography."

In other studies, authors have been concerned about reliability and false-positive results (Wolfer, et al., 2008; Carragee, et al., 2006); some authors have reported high false-positive rates, and others have reported zero false-positive rates. When performed in asymptomatic subjects, the pain provoked by discography cannot be compared to clinical or typical pain; therefore, some studies cannot address true false-positive rates. The patients provoked pain must be similar to the patient's clinical pain for the test to be considered positive.

Additionally, the ability of discography to improve surgical outcomes has yet to be proven (Cohen, et al., 2005) and studies comparing surgical outcomes between patients who have had discography preoperatively with those who have not are few. It is postulated that identification of

RETIRED

Valid for dates of service prior to 11/1/24 only

For dates of service 11/1 and after, see policy:

[EviCore Cigna Commercial Membership | EviCore by Evernorth](#)

a diseased disc as a generator of pain can improve clinical outcomes through better selection of candidates and therapies. In addition it may reduce the likelihood that discs which are not pain generators, are inappropriately treated (Manchikanti, et al., 2009b). When comparing outcomes of fusion procedures lumbar discography is sensitive but lacks specificity (Thiyagarajah, et al., 2021).

The effect of discography on progression of disc degeneration has also gained interest. Carragee et al. (2009) published the results of a prospective matched cohort study evaluating disc degenerative progression over 10 years with (n=50) and without baseline discography (n=52, control). Magnetic resonance imaging was obtained at baseline and at 7-10 years follow-up. The authors noted more frequent and greater degenerative findings, including herniation, end-plate changes, disc grade progression, and annular fissures in the discography group when compared to the matched control group. The authors also noted greater loss of disc height and loss of disc signal in the discography group following annular puncture and injection. In the author's opinion, careful consideration of risk and benefit should be used when recommending procedures involving disc puncture. In 2016, Cueller et al. reported on the comparative incidence of lumbar spine surgery, clinical imaging events, and low back disability events in subjects exposed to discography compared with control subjects over a 10-year follow-up period. At 10-year follow-up, 57 discography subjects and 53 control subjects were able to be included in the final analysis. The authors reported there were 16 lumbar surgeries in the discography group, compared with four in the control group. Medical visits, CT/MRI examinations, work loss, and prolonged back pain episodes were all more frequent in the discography group compared with control subjects (Cueller, et al., 2016).

In an assessment of discography for diagnosis of low back pain, Hayes reviewed seven studies involving sample populations from 36-310 subjects. Comparisons included discography versus no discography prior to surgery (four studies); both groups received discography, then underwent either surgical treatment or nonsurgical management (two studies); pretest/posttest study comparing surgeons' decisions prior to versus after receiving information regarding discography results (one study). Outcomes were measured using Oswestry Disability Index (ODI) scores, quality of life scores (QOL), and change in decision regarding surgery. Follow-up range was six months to six years. Hayes acknowledged the body of evidence is small in size and low in quality and suggests that discography does not lead to improved health outcomes in patients with low back pain being considered for surgery. Two studies suggest discography can lead to serious complications in the long term, including accelerated disc degeneration and increased likelihood of lumbar surgery. According to the authors overall, there is very little evidence supporting the clinical utility of discography (Hayes, 2019).

Evidence in the medical literature supports the use of discography for the following selected conditions (Stretanski and Vu, 2023; Thiyagarajah, et al., 2021; Guyer and Ohnmeiss, 2003):

- for further evaluation of demonstrably abnormal discs when required to assess the extent of the abnormality or correlation of the abnormality with clinical symptoms
- for patients with persistent, severe symptoms in whom other diagnostic tests have failed to reveal clear confirmation of a suspected disc as the source of pain
- for assessment of patients who have failed to respond to previous surgical interventions (i.e., to detect pseudoarthrosis or a symptomatic disc in a posteriorly fused segment and to evaluate for recurrent disc herniation)
- for assessment of discs before fusion
- for assessment of minimally invasive surgical candidates to confirm a contained disc herniation or to investigate dye distribution pattern before chemonucleolysis or other percutaneous procedures

RETIRED

Valid for dates of service prior to 11/1/24 only

For dates of service 11/1 and after, see policy:

[EviCore Cigna Commercial Membership | EviCore by Evernorth](#)

Functional Anesthetic Discography (FAD)

Functional anesthetic discography is a diagnostic procedure that involves injecting an anesthetic agent directly into a spinal disc. Proponents suggest functional anesthetic discography can be used to confirm the presence of injured discs as the source of the patient's low back pain symptoms. According to the manufacturer, functional anesthetic discography is designed to diagnose and potentially treat low back pain caused by degenerative disc disease. Although techniques may vary, during this procedure, under light sedation and x-ray guidance, a small catheter is inserted into the suspected disc and anchored in place with a small balloon. After recovering from light sedation, the patient is asked to engage in physical activity to reproduce pain. Local anesthetic is then injected in the disc believed to be causing the patient's pain. In some cases, administration of intradiscal steroid injection has been proposed in addition to the anesthetic. Reduction in pain is considered diagnostic. If the injection into a specific disc relieves the patient's back pain, the disc can be further evaluated for potential treatment. If the test does not relieve the patient's pain, the physician can investigate other possible causes of pain.

U.S. Food and Drug Administration (FDA): The FAD™ System (originally developed by InnoSpine, Inc., Palo Alto, CA, and later acquired by Kyphon Inc., Sunnyvale, CA) received 510(k) clearance from the FDA in April 2005, as a Class II device. According to the FDA, the intended use of the system is to deliver either a single dose or continuous administration of radiopaque contrast, local anesthetics, and/or saline solution to the intradiscal space. In April 2007 and February 2008, the Discyphor Catheter System (Kyphon, Inc., Sunnyvale, CA), a more recent update to the FAD System, was cleared by the FDA.

Literature Review: Although researchers have investigated the use of functional anesthetic discography for diagnosing discogenic pain, there is insufficient evidence in the published, peer-reviewed scientific literature to support its safety and efficacy at this time. Manchikanti and colleagues (2013b) published an update of the systematic review of the accuracy and utility of lumbar discography in chronic low back pain and reported there was limited evidence supporting FAD or provocation discography with local anesthetic injection.

Contrast Disc Analysis Mapping

The addition of 3D image post-processing, reconstruction and/or mapping of data with markers have been investigated as a method of improving the accuracy and predictive value of discography. However, there is insufficient evidence in the peer-reviewed, published scientific literature to support improved diagnostic utility versus standard provocative discography.

Professional Societies/Organizations

North American Spine Society (NASS): In 2019, NASS issued coverage policy recommendations for the use of discography in the cervical and lumbar spine. Per the recommendations, cervical or lumbar discography is indicated in "the presence of pain and some functional disability for a period of at least 6 months despite conservative therapy. This pain needs to be in a location that could reasonably be caused by the disc (i.e., axial neck or low back, with or without somatic referred pain). In addition, the suspected source of pain identified through other diagnostic imaging testing (e.g., MRI, myelography, CT) needs to be investigated and confirmed and that new or different treatment will be instituted based on the results of the discography. At this time, there are few treatments that would be indicated by a positive discography result; therefore, judicious use of this procedure is indicated" (NASS, 2019).

The NASS guideline outlined the procedural requirements for lumbar discography, as well as consensus guidelines to maximize positive predictive value, minimize false positive tests, and prevent harm:

RETIRED

Valid for dates of service prior to 11/1/24 only

For dates of service 11/1 and after, see policy:

[EviCore Cigna Commercial Membership | EviCore by Evernorth](#)

1. Concordant pain response of $\geq 6/10$ on a modified visual analog scale (VAS) 10 point scale.
2. Volume limit of 3 millimeters (ml).
3. Pressurization of the disc to 15-20 psi above opening pressure, but no greater than 50 pounds per square inch (psi).
4. Adjacent disc(s) provide controls.
 - a. For one control disc:
 - i. Painless response.
 - OR
 - ii. Non-concordant pain that occurs at a pressure >15 psi over opening pressure.
- b. For two adjacent control discs:
 - i. Painless response at both levels.
- OR
- ii. One painless disc AND one disc with non-concordant pain that occurs at a pressure >15 psi over opening pressure.

Regarding cervical spine discography, NASS noted that there is some concern regarding the lack of a standard grading system, as well as the paucity of evidence published. Per NASS, since facet-mediated pain is more prevalent than discogenic pain in the cervical spine, facet-mediated pain should be ruled out with medial branch blocks prior to performing discography in the cervical spine. A disc is considered positive only if stimulation of the target disc reproduces concordant pain with a 7/10 on a visual analogue modified 10 point scale or 70% of most severe pain the patient experiences, and at least one adjacent disc that does not produce pain or produces non-concordant pain with a low volume injection (NASS, 2019).

The 2020 NASS guideline on the diagnosis and treatment of low back (lumbar spine) pain addressed four clinical questions specific to the use of discography. The findings and associated recommendations were as follows (Kreiner, et al., 2020):

- High-level evidence that provocative discography without manometric measurements correlated with pain reproduction in the presence of moderate to severe disc degeneration on MRI/CT discography (Grade of Recommendation: A [Good evidence])
- High-level evidence that provocative discography without manometric pressure measurements correlated with the presence of endplate abnormalities on MRI imaging (Grade of Recommendation: A)
- Conflicting evidence that pressure controlled provocative discography correlates with nuclear transverse relaxation time (T2) signal intensity on magnetic resonance imaging (MRI) in patients with low back pain (Grade of Recommendation: I [Insufficient or conflicting evidence])
- Conflicting evidence that provocative discography without manometric pressure measurements correlates with the presence of a high-intensity zone (HIZ) on MRI imaging (Grade of Recommendation: I)

In both publications discussed above, NASS noted that currently there is not enough high-quality literature available to make a recommendation for or against the use of functional anesthetic discography (Kreiner, et al., 2020; NASS, 2019).

American College of Radiology (ACR): In 2021, the ACR published updated appropriateness criteria for imaging in low back pain. ACR concluded that, although controversial, there was fair evidence supporting the usefulness of lumbar discography for patients with chronic discogenic low back pain, however its use as an initial diagnostic tool in acute uncomplicated low back pain or chronic low back pain without prior management was not supported (ACR, 2021).

RETIRED

Valid for dates of service prior to 11/1/24 only

For dates of service 11/1 and after, see policy:

[EviCore Cigna Commercial Membership | EviCore by Evernorth](#)

The ACR Appropriateness Criteria for Cervical Neck Pain or Cervical Radiculopathy noted that the use of provocative injections in the cervical spine to identify pain is controversial. Plain radiographs MR imaging and CT myelography are more appropriate. Discography is not recommended (ACR, 2018).

American Society of Interventional Pain Physicians (ASIPP): In 2013, ASIPP published updated guidelines on interventional techniques for the management of chronic spinal pain. ASIPP stated that cervical discography is appropriate “only when a treatment is available to test the diagnostic hypothesis of discogenic pain of the cervical spine in individuals who have been properly selected and screened to eliminate other sources of cervical pain”. For thoracic discography, ASIPP noted the evidence was limited and few authors have studied the procedure; however the procedure may be indicated to determine if an intervertebral disc is painful or not, in rare circumstances (based on evidence from two moderate quality studies; there was no recent literature included in the update). Regarding the lumbar spine, ASIPP stated the evidence for diagnostic accuracy of lumbar provocation discography was fair and the evidence for lumbar functional anesthetic discography was limited. Lumbar provocation discography was recommended with appropriate indications in patients with low back pain to prove a diagnostic hypothesis of discogenic pain, specifically after exclusion of other sources of lumbar pain (Manchikanti, et al., 2013a).

American Association of Neurological Surgeons (AANS)/Congress of Neurological Surgeons (CNS): The AANS/CNS Joint Section on Disorders of the Spine and Peripheral Nerves published updated guidelines regarding the performance of fusion procedures for degenerative disease of the lumbar spine (Eck, et al., 2014). According to one of these guidelines, it is recommended that lumbar discography not be used as a stand-alone test on which treatment decisions are based for patients with low back pain. If discography is performed as a diagnostic tool to identify the source of the patient’s low back pain, it is recommended that both a concordant pain response and morphological abnormalities be present at the pathological level prior to initiating any treatment directed at that level. The authors noted that it is possible an association exists between progression of degenerative disc disease and the performance of a provocative discography study; as a result individuals should be counseled regarding potential development prior to undergoing the test.

Medicare Coverage Determinations

	Contractor	Determination Name/Number	Revision Effective Date
NCD		No Determination found	
LCD		No Determination found	

Note: Please review the current Medicare Policy for the most up-to-date information.
(NCD = National Coverage Determination; LCD = Local Coverage Determination)

Coding Information

Notes:

1. This list of codes may not be all-inclusive.
2. Deleted codes and codes which are not effective at the time the service is rendered may not be eligible for reimbursement.

RETIRED

Valid for dates of service prior to 11/1/24 only

For dates of service 11/1 and after, see policy:

[EviCore Cigna Commercial Membership | EviCore by Evernorth](#)

Lumbar Discography

Considered Medically Necessary when criteria in the applicable policy statements listed above are met:

CPT®* Codes	Description
62290	Injection procedure for discography, each level; lumbar
62292	Injection procedure for chemonucleolysis, including discography, intervertebral disc, single or multiple levels, lumbar
72295	Discography, lumbar, radiological supervision and interpretation

Considered Experimental/Investigational/Unproven when used to report cervical or thoracic discography, functional anesthetic discography (FAD) or contrast disc analysis mapping:

CPT®* Codes	Description
62291	Injection procedure for discography, each level; cervical or thoracic
64999	Unlisted procedure, nervous system
72285	Discography, cervical or thoracic, radiological supervision and interpretation
77003	Fluoroscopic guidance and localization of needle or catheter tip for spine or paraspinal diagnostic or therapeutic injection procedures (epidural or subarachnoid) (List separately in addition to code for primary procedure)

***Current Procedural Terminology (CPT®) ©2022 American Medical Association: Chicago, IL.**

References

1. Alamin TF, Kim MJ, Agarwal V. Provocative lumbar discography versus functional anesthetic discography: a comparison of the results of two different diagnostic techniques in 52 patients with chronic low back pain. *Spine J.* 2011 Aug;11(8):756-65.
2. American College of Radiology (ACR). ACR Appropriateness Criteria®: Cervical Neck Pain or Cervical Radiculopathy. Revised 2018. Accessed Oct 3, 2023. Available at URL address: <https://www.acr.org/Clinical-Resources/ACR-Appropriateness-Criteria>
3. American College of Radiology (ACR). ACR Appropriateness Criteria®: Low Back Pain. Revised 2021. Accessed Oct 3, 2023. Available at URL address: <https://www.acr.org/Clinical-Resources/ACR-Appropriateness-Criteria>
4. Boswell MV, Trescot AM, Datta S, Schultz DM, Hansen HC, Abdi S, Sehgal N, Shah RV, Singh V, Benyamin RM, Patel VB, Buenaventura RM, Colson JD, Cordner HJ, Epter RS, Jasper JF, Dunbar EE, Atluri SL, Bowman RC, Deer TR, Swicegood JR, Staats PS, Smith HS, Burton AW, Kloth DS, Giordano J, Manchikanti L; American Society of Interventional Pain Physicians. Interventional techniques: evidence-based practice guidelines in the management of chronic spinal pain. *Pain Physician.* 2007 Jan;10(1):7-111.
5. Brox JI, Nygaard ØP, Holm I, Keller A, Ingebrigtsen T, Reikerås O. Four-year follow-up of surgical versus non-surgical therapy for chronic low back pain. *Ann Rheum Dis.* 2010 Sep;69(9):1643-8.

RETIRED

Valid for dates of service prior to 11/1/24 only

For dates of service 11/1 and after, see policy:

[EviCore Cigna Commercial Membership | EviCore by Evernorth](#)

6. Buenaventura RM, Shah RV, Patel V, Benyamin R, Singh V. Systematic review of discography as a diagnostic test for spinal pain: an update. *Pain Physician*. 2007 Jan;10(1):147-64.
7. Carragee EJ, Alamin TF, Carragee JM. Low-Pressure Positive Discography in Subjects Asymptomatic of Significant Low Back Pain Illness. *Spine*. 2006 Mar;31(5):505-9.
8. Carragee EJ, Don AS, Hurwitz EL, Cuellar JM, Carrino JA, Herzog R. 2009 ISSLS Prize Winner: Does discography cause accelerated progression of degeneration changes in the lumbar disc: a ten-year matched cohort study. *Spine (Phila Pa 1976)*. 2009 Oct 1;34(21):2338-45.
9. Carragee EJ, Hannibal M. Diagnostic evaluation of low back pain. *Orthop Clin N Am*. 2004 Jan;35(1):7-16.
10. Carragee EJ, Lincoln T, Parmar VS, Alamin T. A gold standard evaluation of the "discogenic pain" diagnosis as determined by provocative discography. *Spine*. 2006 Aug 15;31(18):2115-23.
11. Carragee EJ; Tanner CM; Khurana S; Hayward C; Welsh J; Date E; Truong T; Rossi M; Hagle C. The rates of false-positive lumbar discography in select patients without low back symptoms. *Spine*. 2000 Jun;25(11):1373-80; discussion 1381.
12. Carragee EJ, Tanner CM, Yang B, Brito JL, Truong T. False-positive findings on lumbar discography. Reliability of subjective concordance assessment during provocative disc injection. *Spine*. 1999 Dec;24(23):2542-7.
13. Chen Y, Tang T, Erdek MA. Advanced Image-Guided Procedures for Painful Spine. *Neuroimaging Clin N Am*. 2019 Nov;29(4):553-561.
14. Centers for Medicare and Medicaid Services (CMS). Local Coverage Determinations (LCDs) alphabetical index. Accessed Dec 7, 2023. Available at URL address: <https://www.cms.gov/medicare-coverage-database/reports/local-coverage-final-lcds-alphabetical-report.aspx?lcdStatus=all>
15. Centers for Medicare and Medicaid Services (CMS). National Coverage Determinations (NCDs) alphabetical index. Accessed Dec 7, 2023. Available at URL address: <https://www.cms.gov/medicare-coverage-database/reports/national-coverage-ncd-report.aspx?chapter=all&sortBy=title>
16. Cherkin DC, Herman PM. Cognitive and Mind-Body Therapies for Chronic Low Back Pain and Neck Pain: Effectiveness and Value. *JAMA Intern Med*. 2018 Apr 1;178(4):556-557.
17. Chou R, Fu R, Dana T, Pappas M, Hart E, Mauer KM. Interventional Treatments for Acute and Chronic Pain: Systematic Review. Comparative Effectiveness Review No. 247. (Prepared by the Pacific Northwest Evidence-based Practice Center under Contract No. 75Q80120D00006.) AHRQ Publication No. 21-EHC030. Rockville, MD: Agency for Healthcare Research and Quality; Sept 2021.
18. Chou R, Loeser J, Owens D, Rosenquist R, Atlas S, Baisden J, et al. Interventional therapies, surgery, and interdisciplinary rehabilitation for low back pain: an evidence-based

RETIRED

Valid for dates of service prior to 11/1/24 only

For dates of service 11/1 and after, see policy:

[EviCore Cigna Commercial Membership | EviCore by Evernorth](#)

- clinical practice guideline from the American Pain Society. *Spine*. 2009 May 1;34(10):1066-77.
19. Cohen SP, Larkin TM, Barna SA, Palmer WE, Hecht AC, Stojanovic MP. Lumbar discography: a comprehensive review of outcome studies, diagnostic accuracy, and principles. *Reg Anesth Pain Med*. 2005 Mar-Apr;30(2):163-83.
 20. Cuellar JM, Stauff MP, Herzog RJ, Carrino JA, Baker GA, Carragee EJ. Does provocative discography cause clinically important injury to the lumbar intervertebral disc? A 10-year matched cohort study. *Spine J*. 2016 Mar;16(3):273-80.
 21. Derby R, Kim BJ, Lee SH, Chen Y, Seo KS, Aprill C. Comparison of discographic findings in asymptomatic subject discs and the negative discs of chronic LBP patients: can discography distinguish asymptomatic discs among morphologically abnormal discs? *Spine J*. 2005 Jul-Aug;5(4):389-94.
 22. Derby R, Lee SH, Kim BJ, Chen Y, Aprill C, Bogduk N. Pressure-controlled discography in volunteers without low back symptoms. *Pain Med*. 2005 May-Jun;6(3):213-21; discussion 222-4.
 23. Devlin VJ. Chapter 16: Diagnostic and Therapeutic Spinal Injections. In Devlin, ed.: *Spine Secrets*, 3rd ed. Elsevier; 2021:159-168.
 24. Diehn FE, Benson JC, Maus TP. Radiologic Assessment of Patient With Spine Pain. In: Benzon H (ed.). *Practical Management of Pain*. 6th ed. Philadelphia, PA: Elsevier Inc.; 2023. 232-289
 25. Dru AB, Vaziri S, Polifka AJ, Ghogawala Z, Hoh DJ. Indications for Spine Fusion for Axial Pain. In: Steinmetz MP, Berven SH, Benzel EC (eds.). *Benzel's Spine Surgery*. 5th ed. Philadelphia, PA: Elsevier, Inc.; 2022. 1173-1181.e2.
 26. Duthey B. Update on 2004 Background Paper, BP 6.24 Low back pain. Geneva: World Health Organization; 2013.
 27. Eck JC, Sharan A, Resnick DK, Watters WC 3rd, Ghogawala Z, Dailey AT, et al. Guideline update for the performance of fusion procedures for degenerative disease of the lumbar spine. Part 6: discography for patient selection. *Neurosurg Spine*. 2014 Jul;21(1):37-41.
 28. Fairbank J, Frost H, Wilson-MacDonald J, Yu LM, Barker K, Collins R; Spine Stabilisation Trial Group. Randomised controlled trial to compare surgical stabilisation of the lumbar spine with an intensive rehabilitation programme for patients with chronic low back pain: the MRC spine stabilisation trial. *BMJ*. 2005 May 28;330(7502):1233. Erratum in: *BMJ*. 2005 Jun 25;330(7506):1485.
 29. Fang C, Zhang W, Chen L, Li H. The correlation between the high-intensity zone on a T2-weighted MRI and positive outcomes of discography: A meta-analysis. *J Orthop Surg Res*. 2017;12(1):26.
 30. Gardocki RJ, Park AL. Chapter 39: Degenerative Disorders of the Thoracic and Lumbar Spine. In Azar and Beaty, eds: *Campbell's Operative Orthopaedics*, 14th ed. Elsevier; 2021:1719-1801.

RETIRED

Valid for dates of service prior to 11/1/24 only

For dates of service 11/1 and after, see policy:

[EviCore Cigna Commercial Membership | EviCore by Evernorth](#)

31. Grubb SA, Kelly CK. Cervical discography: clinical implications from 12 years of experience. *Spine*. 2000 Jun 1;25(11):1382-9.
32. Gruver C, Guthmiller KB. Provocative Discography. [Updated 2023 May 23]. In: StatPearls [Internet]. Treasure Island (FL): StatPearls Publishing; 2023 Jan-. Accessed Oct 4, 2023. Available at URL address: <https://www.ncbi.nlm.nih.gov/books/NBK470389/>
33. Guyer RD, Ohnmeiss DD; North American Spine Society (NASS). Lumbar discography. *The Spine Journal*. 2003 May-Jun;3(3):11-27.
34. Hayes, Inc. Hayes Medical Technology Directory Report. The Clinical Utility of Lumbar Discography for Assessing Chronic Low Back Pain: Impact on Patient Management and Health Outcomes. Lansdale, PA: Hayes, Inc.; published Sept 7, 2017. Annual review Oct 22, 2018, Nov 12, 2019.
35. Karaarslan N, Yilmaz I, Ozbek H, et al. Are radio-contrast agents commonly used in discography toxic to the intact intervertebral disc tissue cells?. *Basic Clin Pharmacol Toxicol*. 2019;124(2):181-189.
36. Karran EL, Grant AR, Moseley GL. Low back pain and the social determinants of health: a systematic review and narrative synthesis. *Pain*. 2020;161(11):2476-2493.
37. Kreiner DS, Matz P, Bono CM, et al. Guideline summary review: an evidence-based clinical guideline for the diagnosis and treatment of low back pain. *Spine J*. 2020;20(7):998-1024.
38. Laslett M, Aprill CN, McDonald B, Oberg B. Clinical predictors of lumbar provocation discography: a study of clinical predictors of lumbar provocation discography. *Eur Spine J*. 2006 Feb 11;1-12.
39. Madan S, Gundanna M, Harley JM, Boeree NR, Sampson M. Does provocative discography screening of discogenic back pain improve surgical outcome? *J Spinal Disord Tech*. 2002 Jun;15(3):245-51.
40. Manchikanti L, Abdi S, Atluri S, Benyamin RM, Boswell M, Buenaventura RM, et al. An Update of Comprehensive Evidence-Based Guidelines for Interventional Techniques in Chronic Spinal Pain. Part II: Guidance and Recommendations. *Pain Physician* 2013a; 16:S49-S283.
41. Manchikanti L, Benyamin RM, Singh V, Falco FJ, Hameed H, Derby R, et al. An update of the systematic appraisal of the accuracy and utility of lumbar discography in chronic low back pain. *Pain Physician*. 2013b Apr;16(2 Suppl):SE55-95.
42. Manchikanti L, Boswell MV, Singh V, Benyamin RM, Fellows B, Abdi S, Buenaventura RM, Conn A, Datta S, Derby R, Falco FJ, Erhart S, Diwan S, Hayek SM, Helm S, Parr AT, Schultz DM, Smith HS, Wolfer LR, Hirsch JA; ASIPP-IPM. Comprehensive evidence-based guidelines for interventional techniques in the management of chronic spinal pain. *Pain Physician*. 2009c Jul-Aug;12(4):699-802.
43. Manchikanti L, Datta S, Derby R, Wolfer LR, Benyamin RM, Hirsch JA; American Pain Society. A critical review of the American Pain Society clinical practice guidelines for interventional techniques: part 1. Diagnostic interventions. *Pain Physician*. 2010 May-Jun;13(3):E141-74.

RETIRED

Valid for dates of service prior to 11/1/24 only

For dates of service 11/1 and after, see policy:

[EviCore Cigna Commercial Membership | EviCore by Evernorth](#)

44. Manchikanti L, Dunbar EE, Wargo BW, Shah RV, Derby R, Cohen SP. Systematic review of cervical discography as a diagnostic test for chronic spinal pain. *Pain Physician*. 2009a Mar-Apr;12(2):305-21.
45. Manchikanti L, Glaser SE, Wolfer L, Derby R, Cohen SP. Systematic review of lumbar discography as a diagnostic test for chronic low back pain. *Pain Physician*. 2009b May-Jun;12(3):541-59.
46. Manchikanti L, Soin A, Benyamin RM, et al. An Update of the Systematic Appraisal of the Accuracy and Utility of Discography in Chronic Spinal Pain. *Pain Physician*. 2018 Mar;21(2):91-110.
47. Manchikanti L, Staats PS, Singh V, Schultz DM, Vilims BD, Jasper JF, Kloth DS, Trescot AM, Hansen HC, Falasca TD, Racz GB, Deer TR, Burton AW, Helm S, Lou L, Bakhit CE, Dunbar EE, Atluri SL, Calodney AK, Hassenbusch SJ, Feler CA. Evidence-based practice guidelines for interventional techniques in the management of chronic spinal pain. *Pain Physician*. 2003 Jan;6(1):3-81.
48. Margetic P, Pavic R, Stancic MF. Provocative discography screening improves surgical outcome. *Central European Journal of Medicine*. 2013 Oct;125(19-20):600-10.
49. McCormick ZL, Lehman VT, Plastaras CT, et al. Low-Pressure Lumbar Provocation Discography According to Spine Intervention Society/International Association for the Study of Pain Standards Does Not Cause Acceleration of Disc Degeneration in Patients With Symptomatic Low Back Pain: A 7-Year Matched Cohort Study. *Spine (Phila Pa 1976)*. 2019;44(19):E1161-E1168.
50. North American Spine Society (NASS). Discography. Coverage policy recommendations. © 2019 North American Spine Society.
51. North American Spine Society (NASS). Lumbar fusion. Coverage policy recommendations. © 2021 North American Spine Society.
52. Ohnmeiss DD, Guyer RD, Mason SL. The relation between cervical discographic pain responses and radiographic images. *Clin J Pain*. 2000 Mar;16(1):1-5.
53. Onyewu O, Manchikanti L, Falco FJ, et al. An update of the appraisal of the accuracy and utility of cervical discography in chronic neck pain. *Pain Physician*. 2012;15(6):E777-E806.
54. Peh WCG. Provocative discography: current status. *Biomed Imaging Interv J*. 2005;1(1):e2.
55. Poynton AR, Hinman A, Lutz G, Farmer JC. Discography-induced acute lumbar disc herniation. A report of five cases. *J Spinal Disord Tech*. 2005 Apr;18(2):188-92.
56. Resnick DK, Malone DG, Ryken TC. Guidelines for the use of discography for the diagnosis of painful degenerative lumbar disc disease. *Neurosurg Focus*. 2002 Oct;13(2)
57. Santiago K, Cheng J, Jivanelli B, Lutz G. Infections Following Interventional Spine Procedures: A Systematic Review. *Pain Physician*. 2021;24(2):101-116.
58. Shah RV, Everett CR, McKenzie-Brown AM, Sehgal N. Discography as a diagnostic test for spinal pain: A systematic and narrative review. *Pain Physician*. 2005;8:187-209.

RETIRED

Valid for dates of service prior to 11/1/24 only

For dates of service 11/1 and after, see policy:

[EviCore Cigna Commercial Membership | EviCore by Evernorth](#)

59. Shin DA, Kim HI, Jung JH, Shin DG, Lee JO. Diagnostic relevance of pressure-controlled discography. *J Korean Med Sci*. 2006 Oct;21(5):911-6.
60. Smith JS, Sidhu G, Bode K, Gendelberg D, Maltenfort M, Ibrahimi D, Shaffrey CI, Vaccaro AR. Operative and Nonoperative Treatment Approaches for Lumbar Degenerative Disc Disease Have Similar Long-Term Clinical Outcomes Among Patients with Positive Discography. *World Neurosurg*. 2013 Sep 15. pii: S1878-8750(13)01111-X.
61. Staartjes VE, Vergroesen PA, Zeilstra DJ, Schröder ML. Identifying subsets of patients with single-level degenerative disc disease for lumbar fusion: the value of prognostic tests in surgical decision making. *Spine J*. 2018 Apr;18(4):558-566.
62. Stretanski MF, Vu L. Fluoroscopy Discography Assessment, Protocols, and Interpretation. [Updated 2023 Jul 10]. In: StatPearls [Internet]. Treasure Island (FL): StatPearls Publishing; 2023 Jan-. Accessed Oct 4, 2023. Available at URL address: <https://www.ncbi.nlm.nih.gov/books/NBK572119/>
63. Thiyagarajah AR, Hord ED, Vallejo R. Discography. Medscape. Updated Aug 10, 2021. Accessed Oct 4, 2023. Available at URL address: <https://emedicine.medscape.com/article/1145703-overview>
64. U.S. Food and Drug Administration (FDA). Discyphor™ Catheter System. 510(k) Summary. K063071. Apr 13, 2007. Accessed Oct 4, 2023. Available at URL address: <https://www.accessdata.fda.gov/scripts/cdrh/cfdocs/cfPMN/pmn.cfm>
65. U.S. Food and Drug Administration (FDA). Discyphor™ Catheter System. 510(k) Summary. K073516. Feb 21, 2008. Accessed Oct 4, 2023. Available at URL address: <https://www.accessdata.fda.gov/scripts/cdrh/cfdocs/cfPMN/pmn.cfm>
66. U.S. Food and Drug Administration (FDA). Functional Anaesthetic Discography System. 510(k) Summary. K043500. Apr 15, 2005. Accessed Oct 4, 2023. Available at URL address: <https://www.accessdata.fda.gov/scripts/cdrh/cfdocs/cfPMN/pmn.cfm>
67. Wieser ES, Wang JC. Surgery for neck pain. *Neurosurgery*. 2007 Jan;60(1 Suppl 1):S51-6.
68. Willems PC, Elmans L, Anderson PG, van der Schaaf DB, de Kleuver M. Provocative discography and lumbar fusion: is preoperative assessment of adjacent discs useful? *Spine*. 2007 May 1;32(10):1094-9; discussion 1100.
69. Wolfer LR, Derby R, Lee JE, Lee SH. Systematic review of lumbar provocation discography in asymptomatic subjects with a meta-analysis of false-positive rates. *Pain Physician*. 2008 Jul-Aug;11(4):513-38.
70. Wood KB, Schellhas KP, Garvey TA, Aeppli D. Thoracic discography in healthy individuals. A controlled prospective study of magnetic resonance imaging and discography in asymptomatic and symptomatic individuals. *Spine (Phila Pa 1976)*. 1999;24(15):1548-1555.
71. Wu PH, Kim HS, Jang IT. Intervertebral Disc Diseases PART 2: A Review of the Current Diagnostic and Treatment Strategies for Intervertebral Disc Disease. *Int J Mol Sci*. 2020b;21(6):2135. Published 2020 Mar 20.

RETIRED

Valid for dates of service prior to 11/1/24 only

For dates of service 11/1 and after, see policy:

[EviCore Cigna Commercial Membership | EviCore by Evernorth](#)

72. Wu A, March L, Zheng X, et al. Global low back pain prevalence and years lived with disability from 1990 to 2017: estimates from the Global Burden of Disease Study 2017. *Ann Transl Med.* 2020a;8(6):299.
73. Xi MA, Tong HC, Fahim DK, Perez-Cruet M. Using Provocative Discography and Computed Tomography to Select Patients with Refractory Discogenic Low Back Pain for Lumbar Fusion Surgery. *Cureus.* 2016 Feb 27;8(2):e514.
74. Zheng Y, Liew SM, Simmons ED. Value of magnetic resonance imaging and discography in determining the level of cervical discectomy and fusion. *Spine.* 2004 Oct;29(19):2140-5, discussion 2146.
75. Zhou Y, Abdi S. Diagnosis and minimally invasive treatment of lumbar discogenic pain--a review of the literature. *Clin J Pain.* 2006 Jun;22(5):468-81.

Revision Details

Type of Revision	Summary of Changes	Date
Annual review	<ul style="list-style-type: none">• No policy statement changes.	11/15/2023

"Cigna Companies" refers to operating subsidiaries of The Cigna Group. All products and services are provided exclusively by or through such operating subsidiaries, including Cigna Health and Life Insurance Company, Connecticut General Life Insurance Company, Evernorth Behavioral Health, Inc., Cigna Health Management, Inc., and HMO or service company subsidiaries of The Cigna Group. © 2023 The Cigna Group.