



## Drug Coverage Policy

Effective Date .....8/1/2024

Coverage Policy Number.....IP0152

Policy Title.....Doptelet

# Thrombocytopenia – Doptelet

- Doptelet® (avatrombopag tablets - Dova/AkaRx)

### INSTRUCTIONS FOR USE

The following Coverage Policy applies to health benefit plans administered by Cigna Companies. Certain Cigna Companies and/or lines of business only provide utilization review services to clients and do not make coverage determinations. References to standard benefit plan language and coverage determinations do not apply to those clients. Coverage Policies are intended to provide guidance in interpreting certain standard benefit plans administered by Cigna Companies. Please note, the terms of a customer’s particular benefit plan document [Group Service Agreement, Evidence of Coverage, Certificate of Coverage, Summary Plan Description (SPD) or similar plan document] may differ significantly from the standard benefit plans upon which these Coverage Policies are based. For example, a customer’s benefit plan document may contain a specific exclusion related to a topic addressed in a Coverage Policy. In the event of a conflict, a customer’s benefit plan document always supersedes the information in the Coverage Policies. In the absence of a controlling federal or state coverage mandate, benefits are ultimately determined by the terms of the applicable benefit plan document. Coverage determinations in each specific instance require consideration of 1) the terms of the applicable benefit plan document in effect on the date of service; 2) any applicable laws/regulations; 3) any relevant collateral source materials including Coverage Policies and; 4) the specific facts of the particular situation. Each coverage request should be reviewed on its own merits. Medical directors are expected to exercise clinical judgment and have discretion in making individual coverage determinations. Coverage Policies relate exclusively to the administration of health benefit plans. Coverage Policies are not recommendations for treatment and should never be used as treatment guidelines. In certain markets, delegated vendor guidelines may be used to support medical necessity and other coverage determinations.

## Cigna Healthcare Coverage Policy

### OVERVIEW

Doptelet, a thrombopoietin receptor agonist, is indicated for the following uses:<sup>1</sup>

- **Immune thrombocytopenia (ITP)**, chronic, for treatment in adults who have had an insufficient response to a previous treatment.
- **Thrombocytopenia**, as treatment in adults with **chronic liver disease** who are scheduled to undergo a procedure.

For chronic ITP, Doptelet should be discontinued if the platelet count does not increase to  $\geq 50 \times 10^9/L$  within 4 weeks at the maximum dose of 40 mg once daily. The safety and efficacy of Doptelet have not been established in pediatric patients. For chronic liver disease in patients

undergoing a procedure, Doptelet is given as a 5-day course beginning 10 to 13 days before the scheduled procedure. In general, patients in the pivotal studies had a platelet count  $< 50 \times 10^9/L$ .

### Guidelines

In 2019, the American Society of Hematology updated guidelines for ITP.<sup>4</sup> Doptelet is not addressed specifically, but there are several recommendations. For adults with ITP for at least 3 months who are corticosteroid-dependent or unresponsive to corticosteroid, a thrombopoietin receptor agonist (either Promacta<sup>®</sup> [eltrombopag tablets and oral suspension] or Nplate<sup>®</sup> [romiplostim subcutaneous injection]) or a splenectomy are recommended. In children with newly diagnosed ITP who have non-life-threatening mucosal bleeding, corticosteroids are recommended. For children who have non-life-threatening mucosal bleeding and did not respond to first-line treatment, thrombopoietin receptor agonists are recommended. Other treatment options in children and adults include intravenous immunoglobulin, anti-D immunoglobulin, and rituximab.

## Medical Necessity Criteria

**Doptelet is considered medically necessary when ONE of the following is met (1 or 2):**

### FDA-Approved Indications

- 1. Chronic Immune Thrombocytopenia.** Approve if the patient meets ALL of the following (A or B):
  - A) Initial Therapy.** Approve for 3 months if the patient meets ALL of the following (i, ii, iii, and iv):
    - i. Patient is  $\geq 18$  years of age; AND
    - ii. Patient meets ONE of the following (a or b):
      - a) Patient has a platelet count  $< 30 \times 10^9/L$  ( $< 30,000/mcL$ ); OR
      - b) Patient meets BOTH of the following [(1) and (2)]:
        - (1) Patient has a platelet count  $< 50 \times 10^9/L$  ( $< 50,000/mcL$ ); AND
        - (2) According to the prescriber, the patient is at an increased risk of bleeding; AND
    - iii. Patient meets ONE of the following (a or b):
      - a) Patient has tried at least ONE other therapy; OR  
Note: Examples of therapies are systemic corticosteroids, intravenous immunoglobulin, anti-D immunoglobulin, Promacta (eltrombopag tablets and oral suspension), Nplate (romiplostim subcutaneous injection), Tavalisse (fostamatinib tablets), and rituximab.
      - b) Patient has undergone splenectomy; AND
    - iv. The medication is prescribed by or in consultation with a hematologist; OR
  - B) Patient is Currently Receiving Doptelet.** Approve for 1 year if the patient meets BOTH of the following (i and ii):
    - i. According to the prescriber, the patient demonstrates a beneficial clinical response; AND  
Note: A beneficial response can include increased platelet counts, maintenance of platelet counts, and/or a decreased frequency of bleeding episodes.
    - ii. Patient remains at risk for bleeding complications.
- 2. Thrombocytopenia in a Patient with Chronic Liver Disease.** Approve for 5 days if the patient meets ALL of the following (A, B, and C):
  - A) Patient is  $\geq 18$  years of age; AND**
  - B) Patient has a current platelet count  $< 50 \times 10^9/L$  ( $< 50,000/mcL$ ); AND**
  - C) Patient is scheduled to undergo a procedure within 10 to 13 days after starting Doptelet therapy.**

When coverage is available and medically necessary, the dosage, frequency, duration of therapy, and site of care should be reasonable, clinically appropriate, and supported by evidence-based literature and adjusted based upon severity, alternative available treatments, and previous response to therapy.

Receipt of sample product does not satisfy any criteria requirements for coverage.

## Conditions Not Covered

Any other use is considered experimental, investigational, or unproven (criteria will be updated as new published data are available).

## References

1. Doptelet® tablets [prescribing information]. Durham, NC: AkaRx/Dova; July 2021.
2. Neunert C, Terrell DR, Arnold DM, et al. American Society of Hematology 2019 guidelines for immune thrombocytopenia. *Blood Adv.* 2019;3(23):3829-3866.

## Revision Details

Type of Revision	Summary of Changes	Date
Annual Revision	<p><b>Updated</b> policy title to Thrombocytopenia – Doptelet; previously was Avatrombopag.</p> <p><b>Chronic Immune Thrombocytopenia:</b> Updated pre-requisite therapy requirement from “Individual has had an inadequate response to ONE of the following OR Individual has a contraindication or is intolerant to ALL of the following: Systemic corticosteroids, Intravenous immunoglobulin, Anti-D immunoglobulin, Promacta, Nplate, Tavalisse, Rituximab” to now be “Patient has tried at least ONE other therapy” with examples relocated to a Note.</p>	8/1/2024

The policy effective date is in force until updated or retired.

---

“Cigna Companies” refers to operating subsidiaries of The Cigna Group. All products and services are provided exclusively by or through such operating subsidiaries, including Cigna Health and Life Insurance Company, Connecticut General Life Insurance Company, Evernorth Behavioral Health, Inc., Cigna Health Management, Inc., and HMO or service company subsidiaries of The Cigna Group. © 2024 The Cigna Group.