



Drug Coverage Policy

Effective Date.....7/1/2024
Coverage Policy Number.....IP0280
Policy Title.....Saphnelo

Lupus - Saphnelo

- Saphnelo® (anifrolumab-fnia intravenous infusion – AstraZeneca)

INSTRUCTIONS FOR USE

The following Coverage Policy applies to health benefit plans administered by Cigna Companies. Certain Cigna Companies and/or lines of business only provide utilization review services to clients and do not make coverage determinations. References to standard benefit plan language and coverage determinations do not apply to those clients. Coverage Policies are intended to provide guidance in interpreting certain standard benefit plans administered by Cigna Companies. Please note, the terms of a customer’s particular benefit plan document [Group Service Agreement, Evidence of Coverage, Certificate of Coverage, Summary Plan Description (SPD) or similar plan document] may differ significantly from the standard benefit plans upon which these Coverage Policies are based. For example, a customer’s benefit plan document may contain a specific exclusion related to a topic addressed in a Coverage Policy. In the event of a conflict, a customer’s benefit plan document always supersedes the information in the Coverage Policies. In the absence of a controlling federal or state coverage mandate, benefits are ultimately determined by the terms of the applicable benefit plan document. Coverage determinations in each specific instance require consideration of 1) the terms of the applicable benefit plan document in effect on the date of service; 2) any applicable laws/regulations; 3) any relevant collateral source materials including Coverage Policies and; 4) the specific facts of the particular situation. Each coverage request should be reviewed on its own merits. Medical directors are expected to exercise clinical judgment and have discretion in making individual coverage determinations. Coverage Policies relate exclusively to the administration of health benefit plans. Coverage Policies are not recommendations for treatment and should never be used as treatment guidelines. In certain markets, delegated vendor guidelines may be used to support medical necessity and other coverage determinations.

Cigna Healthcare Coverage Policy

OVERVIEW

Saphnelo, a type 1 interferon (IFN) receptor antagonist, is indicated for the treatment of moderate to severe **systemic lupus erythematosus (SLE)** in adults who are receiving standard therapy.¹

Saphnelo efficacy has not been evaluated and is not recommended in patients with severe active lupus nephritis or severe active central nervous system lupus.

Guidelines

European League Against Rheumatism (EULAR) guidelines for SLE (2023) recommend hydroxychloroquine for all patients, unless contraindicated.² Depending on the type and severity of organ involvement, glucocorticoids can be used but dosing should be minimized or withdrawn. Methotrexate, azathioprine, mycophenolate, and/or biologic agents (Benlysta® [belimumab intravenous or subcutaneous infusion], Saphnelo) should be considered in patients who do not respond to hydroxychloroquine ± glucocorticoids. EULAR also states biologic agents (Benlysta, Saphnelo) should be considered as second-line therapy for the treatment of active skin disease. Patient with active proliferative lupus nephritis should also consider combination therapy with biologic agents (Benlysta, Lupkynis™ [voclosporin capsules]). In general, the pharmacological interventions are directed by patient characteristics and the type/severity of organ involvement.

Medical Necessity Criteria

Saphnelo is considered medically necessary when the following criteria are met:

FDA-Approved Indication

1. Systemic Lupus Erythematosus. Approve for the duration noted if the patient meets ONE of the following (A or B):

- A) Initial Therapy. Approve for 6 months if the patient meets ALL of the following (i, ii, iii, iv and v):
- i. Patient is ≥ 18 years of age; AND
 - ii. Patient has autoantibody-positive SLE, defined as positive for at least one of the following: antinuclear antibodies (ANA), anti-double-stranded DNA (anti-dsDNA) antibodies, anti-Smith (anti-Sm) antibodies; AND
Note: Not all patients with SLE are positive for anti-dsDNA, but most will be positive for ANA.
 - iii. Patient meets ONE of the following (a or b):
 - a) The medication is being used concurrently with at least one other standard therapy; OR
Note: Examples of standard therapies include an antimalarial (e.g., hydroxychloroquine), systemic corticosteroid (e.g., prednisone), and other immunosuppressants (e.g., azathioprine, mycophenolate mofetil, methotrexate).
 - b) Patient is determined to be intolerant to standard therapy due to a significant toxicity, as determined by the prescriber; AND
 - iv. The medication is prescribed by or in consultation with a rheumatologist, clinical immunologist, nephrologist, neurologist, or dermatologist.
 - v. Preferred product criteria is met for the product(s) as listed in the below table(s)

Employer Plans:

Product	Criteria
Saphnelo (anifrolumab-fnia)	ONE of the following: <ol style="list-style-type: none"> 1. Failure, contraindication, or intolerance to Benlysta (belimumab) [may require prior authorization] intravenous infusion or subcutaneous injection. 2. Depression or suicidality, according to the prescriber. 3. Treatment with Saphnelo has been started.

- B) Patient is Currently Receiving Saphnelo. Approve for 1 year if the patient meets ALL of the following (i, ii, and iii):
- i. Patient meets ONE of the following (a or b):

- a) The medication is being used concurrently with at least one other standard therapy;
OR
Note: Examples of standard therapies include an antimalarial (e.g., hydroxychloroquine), systemic corticosteroid (e.g., prednisone), and other immunosuppressants (e.g., azathioprine, mycophenolate mofetil, methotrexate).
- b) Patient is determined to be intolerant to standard therapy due to a significant toxicity, as determined by the prescriber; AND
- ii. Patient responded to Saphnelo, as determined by the prescriber; AND
Note: Examples of a response include reduction in flares, reduction in corticosteroid dose, decrease of anti-dsDNA titer, improvement in complement levels (i.e., C3, C4), or improvement in specific organ dysfunction (e.g., musculoskeletal, blood, hematologic, vascular, others).
- iii. The medication is prescribed by or in consultation with a rheumatologist, clinical immunologist, nephrologist, neurologist, or dermatologist.

Dosing. Approve 300 mg given as an intravenous infusion administered not more frequently than once every 4 weeks.

When coverage is available and medically necessary, the dosage, frequency, duration of therapy, and site of care should be reasonable, clinically appropriate, and supported by evidence-based literature and adjusted based upon severity, alternative available treatments, and previous response to therapy.

Receipt of sample product does not satisfy any criteria requirements for coverage.

Conditions Not Covered

Any other use is considered experimental, investigational, or unproven, including the following (this list may not be all inclusive; criteria will be updated as new published data are available): Coverage of Saphnelo is not recommended in the following situations:

- 1. Concurrent Use with Other Biologics.** Saphnelo has not been studied and is not recommended in combination with other biologics (e.g., Benlysta [belimumab intravenous infusion or subcutaneous injection], rituximab).¹ Safety and efficacy have not been established with these combinations. See [APPENDIX](#) for examples of other biologics that should not be taken in combination with Saphnelo.

Coding Information

- 1) This list of codes may not be all-inclusive.
- 2) Deleted codes and codes which are not effective at the time the service is rendered may not be eligible for reimbursement.

Considered Medically Necessary when criteria in the applicable policy statements listed above are met:

HCPCS Codes	Description
J0491	Injection, anifrolumab-fnia, 1 mg

References

1. Saphnelo® injection, for intravenous use [prescribing information]. Wilmington DE: AstraZeneca; September 2022.
2. Fanouriakis A, Kostopoulou M, Andersen J, et al. EULAR recommendations for the management of systemic lupus erythematosus: 2023 update. *Ann Rheum Dis.* 2024;83(1):15-29.

Revision Details

Type of Revision	Summary of Changes	Date
Annual Revision	<p>Updated coverage policy title from <i>Anifrolumab-fnia</i> to <i>Lupus Saphnelo</i>.</p> <p>No criteria changes</p>	6/15/2024

The policy effective date is in force until updated or retired.

APPENDIX

	Mechanism of Action	Examples of Inflammatory Indications*
Biologics		
Saphnelo™ (anifrolumab-fnia IV infusion)	IFN receptor antagonist	SLE
Adalimumab SC Products (Humira®, biosimilars)	Inhibition of TNF	AS, CD, JIA, PsO, PsA, RA, UC
Cimzia® (certolizumab pegol SC injection)	Inhibition of TNF	AS, CD, nr-axSpA, PsO, PsA, RA
Etanercept SC Products (Enbrel®, biosimilars)	Inhibition of TNF	AS, JIA, PsO, PsA
Infliximab IV Products (Remicade®, biosimilars)	Inhibition of TNF	AS, CD, PsO, PsA, RA, UC
Simponi®, Simponi® Aria™ (golimumab SC injection, golimumab IV infusion)	Inhibition of TNF	SC formulation: AS, PsA, RA, UC
		IV formulation: AS, PJIA, PsA, RA
Actemra® (tocilizumab IV infusion, tocilizumab SC injection)	Inhibition of IL-6	SC formulation: PJIA, RA, SJIA
		IV formulation: PJIA, RA, SJIA
Kevzara® (sarilumab SC injection)	Inhibition of IL-6	RA
Orencia® (abatacept IV infusion, abatacept SC injection)	T-cell costimulation modulator	SC formulation: JIA, PsA, RA
		IV formulation: JIA, PsA, RA
Rituximab IV Products (Rituxan®, biosimilars)	CD20-directed cytolytic antibody	RA
Kineret® (anakinra SC injection)	Inhibition of IL-1	JIA^, RA
Stelara® (ustekinumab SC injection, ustekinumab IV infusion)	Inhibition of IL-12/23	SC formulation: CD, PsO, PsA, UC
		IV formulation: CD, UC

Siliq™ (brodalumab SC injection)	Inhibition of IL-17	PsO
Cosentyx® (secukinumab SC injection)	Inhibition of IL-17A	AS, ERA, nr-axSpA, PsO, PsA
Taltz® (ixekizumab SC injection)	Inhibition of IL-17A	AS, nr-axSpA, PsO, PsA
Ilumya™ (tildrakizumab-asmn SC injection)	Inhibition of IL-23	PsO
Skyrizi® (risankizumab-rzaa SC injection, risankizumab-rzaa IV infusion)	Inhibition of IL-23	SC formulation: CD, PsA, PsO
		IV formulation: CD
Tremfya™ (guselkumab SC injection)	Inhibition of IL-23	PsO, PsA
Entyvio™ (vedolizumab IV infusion)	Integrin receptor antagonist	CD, UC

* Not an all-inclusive list of indications (e.g., oncology indications and rare inflammatory conditions are not listed). Refer to the prescribing information for the respective agent for FDA-approved indications; IFN – Interferon; SLE – Systemic lupus erythematosus; SC – Subcutaneous; TNF – Tumor necrosis factor; AS – Ankylosing spondylitis; CD – Crohn’s disease; JIA – Juvenile idiopathic arthritis; PsO – Plaque psoriasis; PsA – Psoriatic arthritis; RA – Rheumatoid arthritis; UC – Ulcerative colitis; nr-axSpA – Non-radiographic axial spondyloarthritis; IV – Intravenous, PJIA – Polyarticular juvenile idiopathic arthritis; IL – Interleukin; SJIA – Systemic juvenile idiopathic arthritis; ^ Off-label use of Kineret in JIA supported in guidelines; ERA – Enthesitis-related arthritis.

“Cigna Companies” refers to operating subsidiaries of The Cigna Group. All products and services are provided exclusively by or through such operating subsidiaries, including Cigna Health and Life Insurance Company, Connecticut General Life Insurance Company, Evernorth Behavioral Health, Inc., Cigna Health Management, Inc., and HMO or service company subsidiaries of The Cigna Group. © 2024 The Cigna Group.