



## Drug Coverage Policy

Effective Date.....11/01/2024  
Coverage Policy Number.....IP0691  
Policy Title.....Velsipity Prior  
Authorization Policy

# Inflammatory Conditions – Velsipity Prior Authorization Policy

- Velsipity® (etrasimod tablets – Pfizer)

### **INSTRUCTIONS FOR USE**

*The following Coverage Policy applies to health benefit plans administered by Cigna Companies. Certain Cigna Companies and/or lines of business only provide utilization review services to clients and do not make coverage determinations. References to standard benefit plan language and coverage determinations do not apply to those clients. Coverage Policies are intended to provide guidance in interpreting certain standard benefit plans administered by Cigna Companies. Please note, the terms of a customer's particular benefit plan document [Group Service Agreement, Evidence of Coverage, Certificate of Coverage, Summary Plan Description (SPD) or similar plan document] may differ significantly from the standard benefit plans upon which these Coverage Policies are based. For example, a customer's benefit plan document may contain a specific exclusion related to a topic addressed in a Coverage Policy. In the event of a conflict, a customer's benefit plan document always supersedes the information in the Coverage Policies. In the absence of a controlling federal or state coverage mandate, benefits are ultimately determined by the terms of the applicable benefit plan document. Coverage determinations in each specific instance require consideration of 1) the terms of the applicable benefit plan document in effect on the date of service; 2) any applicable laws/regulations; 3) any relevant collateral source materials including Coverage Policies and; 4) the specific facts of the particular situation. Each coverage request should be reviewed on its own merits. Medical directors are expected to exercise clinical judgment and have discretion in making individual coverage determinations. Coverage Policies relate exclusively to the administration of health benefit plans. Coverage Policies are not recommendations for treatment and should never be used as treatment guidelines. In certain markets, delegated vendor guidelines may be used to support medical necessity and other coverage determinations.*

## Cigna Healthcare Coverage Policy

### **Overview**

Velsipity, a sphingosine 1-phosphate receptor modulator, is indicated for the treatment of **ulcerative colitis** (UC), in adults with moderately to severely active disease.<sup>1</sup>

### **Guidelines/Clinical Efficacy**

Velsipity is not currently addressed in UC guidelines. The American Gastroenterological Association (2020) and the American College of Gastroenterology (2019) have clinical practice guidelines on the management of moderate to severe UC and make recommendations for induction and maintenance of remission in adults.<sup>2,3</sup> Both endorse the use of biologic agents and give specific patient circumstances in the selection for induction and maintenance therapies. Pivotal trials for Velsipity included adults with moderately to severely active UC who had an inadequate response or were intolerant to any of the following agents: oral aminosalicylates, corticosteroids, immunomodulators (e.g., 6-mercaptopurine and azathioprine), or a biologic (e.g., tumor necrosis factor inhibitor, Entyvio® [vedolizumab injection], or a Janus kinase inhibitor (e.g., Xeljanz® [tofacitinib tablets]).<sup>1</sup>

## Medical Necessity Criteria

### Policy Statement

Prior Authorization is recommended for benefit coverage of Velsipity. All approvals are provided for the duration noted below. In cases where the approval is authorized in months, 1 month is equal to 30 days. Because of the specialized skills required for evaluation and diagnosis of patients treated with Velsipity as well as the monitoring required for adverse events and long-term efficacy, approval requires Velsipity to be prescribed by or in consultation with a physician who specializes in the condition being treated.

**Velsipity is considered medically necessary when the following are met:**

### FDA-Approved Indications

- 1. Ulcerative Colitis.** Approve for the duration noted if the patient meets ONE of the following (A or B):
  - A) Initial Therapy.** Approve for 6 months if the patient meets ALL of the following (i, ii, and iii):
    - i.** Patient is  $\geq$  18 years of age; AND
    - ii.** Patient has had a trial of ONE systemic agent for ulcerative colitis; AND  
Note: Examples of systemic agents for ulcerative colitis include 6-mercaptopurine, azathioprine, cyclosporine, tacrolimus, or a corticosteroid such as prednisone, methylprednisolone. A trial of one biologic also counts as a trial of one systemic agent for ulcerative colitis. Refer to the [Appendix](#) for examples of biologics used for ulcerative colitis.
    - iii.** The medication is prescribed by or in consultation with a gastroenterologist.
  - B) Patient is Currently Receiving Velsipity.** Approve for 1 year if the patient meets BOTH of the following (i and ii):
    - i.** Patient has been established on therapy for at least 6 months; AND  
Note: A patient who has received < 6 months of therapy or who is restarting therapy is reviewed under criterion A (Initial Therapy).
    - ii.** Patient meets at least one of the following (a or b):
      - a)** When assessed by at least one objective measure, patient experienced a beneficial clinical response from baseline (prior to initiating the requested drug); OR  
Note: Examples of assessment for inflammatory response include fecal markers (e.g., fecal calprotectin), serum markers (e.g., C-reactive protein), endoscopic assessment, and/or reduced dose of corticosteroids.
      - b)** Compared with baseline (prior to initiating Velsipity), patient experienced an improvement in at least one symptom, such as decreased pain, fatigue, stool frequency, and/or decreased rectal bleeding.

When coverage is available and medically necessary, the dosage, frequency, duration of therapy, and site of care should be reasonable, clinically appropriate, and supported by evidence-based literature and adjusted based upon severity, alternative available treatments, and previous response to therapy.

Receipt of sample product does not satisfy any criteria requirements for coverage.

## Conditions Not Covered

Any other use is considered experimental, investigational, or unproven, including the following (this list may not be all inclusive; criteria will be updated as new published data are available):

- 1. Concurrent Use with a Biologic or with a Targeted Synthetic Oral Small Molecule Drug.** This medication should not be administered in combination with another biologic or with a targeted synthetic oral small molecule drug used for an inflammatory condition (see [Appendix](#) for examples). Combination therapy is generally not recommended due to a potentially higher rate of adverse events and lack of controlled clinical data supporting additive efficacy.
- 2. Concurrent Use with Other Potent Immunosuppressants.** In pivotal trials, patients who received Velsipity were not to receive concomitant treatment with non-corticosteroid immunosuppressive or immune-modulating therapies used for the treatment of ulcerative colitis. Combination therapy is generally not recommended due to a potential for a higher rate of adverse effects with combinations and lack of controlled clinical data supporting additive efficacy.<sup>1</sup>  
Note: Examples include 6-mercaptopurine, azathioprine, cyclosporine, and methotrexate.

## References

1. Velsipity® tablets [prescribing information]. New York, NY: Pfizer; October 2023.
2. Feuerstein JD, Isaac s KL, Schneider Y, et al. AGA clinical practice guidelines on the management of moderate to severe ulcerative colitis. *Gastroenterology*. 2020;158:1450-1461.
3. Rubin DT, Ananthkrishnan AN, Siegel CA, et al. American College of Gastroenterology clinical guideline: ulcerative colitis in adults. *Am J Gastroenterol*. 2019;114:384-413.

### APPENDIX

	Mechanism of Action	Examples of Indications*
<b>Biologics</b>		
<b>Adalimumab SC Products</b> (Humira®, biosimilars)	Inhibition of TNF	AS, CD, JIA, PsO, PsA, RA, UC
<b>Cimzia®</b> (certolizumab pegol SC injection)	Inhibition of TNF	AS, CD, nr-axSpA, PsO, PsA, RA
<b>Etanercept SC Products</b> (Enbrel®, biosimilars)	Inhibition of TNF	AS, JIA, PsO, PsA, RA
<b>Infliximab IV Products</b> (Remicade®, biosimilars)	Inhibition of TNF	AS, CD, PsO, PsA, RA, UC
<b>Zymfentra®</b> (infliximab-dyyb SC injection)	Inhibition of TNF	CD, UC
<b>Simponi®, Simponi Aria®</b> (golimumab SC injection, golimumab IV infusion)	Inhibition of TNF	SC formulation: AS, PsA, RA, UC
		IV formulation: AS, PJIA, PsA, RA
<b>Tocilizumab Products</b> (Actemra® IV, biosimilar; Actemra SC, biosimilar)	Inhibition of IL-6	SC formulation: PJIA, RA, SJIA

		IV formulation: PJIA, RA, SJIA
<b>Kevzara</b> <sup>®</sup> (sarilumab SC injection)	Inhibition of IL-6	RA
<b>Orencia</b> <sup>®</sup> (abatacept IV infusion, abatacept SC injection)	T-cell costimulation modulator	SC formulation: JIA, PSA, RA IV formulation: JIA, PsA, RA
<b>Rituximab IV Products</b> (Rituxan <sup>®</sup> , biosimilars)	CD20-directed cytolytic antibody	RA
<b>Kineret</b> <sup>®</sup> (anakinra SC injection)	Inhibition of IL-1	JIA <sup>^</sup> , RA
<b>Omvo</b> <sup>®</sup> (mirikizumab IV infusion, SC injection)	Inhibition of IL-23	UC
<b>Stelara</b> <sup>®</sup> (ustekinumab SC injection, ustekinumab IV infusion)	Inhibition of IL-12/23	SC formulation: CD, PsO, PsA, UC IV formulation: CD, UC
<b>Siliq</b> <sup>®</sup> (brodalumab SC injection)	Inhibition of IL-17	PsO
<b>Cosentyx</b> <sup>®</sup> (secukinumab SC injection; secukinumab IV infusion)	Inhibition of IL-17A	SC formulation: AS, ERA, nr-axSpA, PsO, PsA IV formulation: AS, nr-axSpA, PsA
<b>Taltz</b> <sup>®</sup> (ixekizumab SC injection)	Inhibition of IL-17A	AS, nr-axSpA, PsO, PsA
<b>Bimzelx</b> <sup>®</sup> (bimekizumab-bkzx SC injection)	Inhibition of IL-17A/17F	PsO
<b>Ilumya</b> <sup>®</sup> (tildrakizumab-asmn SC injection)	Inhibition of IL-23	PsO
<b>Skyrizi</b> <sup>®</sup> (risankizumab-rzaa SC injection, risankizumab-rzaa IV infusion)	Inhibition of IL-23	SC formulation: CD, PSA, PsO, UC IV formulation: CD, UC
<b>Tremfya</b> <sup>®</sup> (guselkumab SC injection, guselkumab IV infusion)	Inhibition of IL-23	SC formulation: PsA, PsO, UC IV formulation: UC
<b>Entyvio</b> <sup>®</sup> (vedolizumab IV infusion, vedolizumab SC injection)	Integrin receptor antagonist	CD, UC
<b>Oral Therapies/Targeted Synthetic Oral Small Molecule Drugs</b>		
<b>Otezla</b> <sup>®</sup> (apremilast tablets)	Inhibition of PDE4	PsO, PsA
<b>Cibinqo</b> <sup>™</sup> (abrocitinib tablets)	Inhibition of JAK pathways	AD
<b>Olumiant</b> <sup>®</sup> (baricitinib tablets)	Inhibition of JAK pathways	RA, AA
<b>Litfulo</b> <sup>®</sup> (ritlecitinib capsules)	Inhibition of JAK pathways	AA
<b>Leqselvi</b> <sup>®</sup> (deuruxolitinib tablets)	Inhibition of JAK pathways	AA
<b>Rinvoq</b> <sup>®</sup> (upadacitinib extended-release tablets)	Inhibition of JAK pathways	AD, AS, nr-axSpA, RA, PsA, UC
<b>Rinvoq</b> <sup>®</sup> LQ (upadacitinib oral solution)	Inhibition of JAK pathways	PsA, PJIA
<b>Sotyktu</b> <sup>®</sup> (deucravacitinib tablets)	Inhibition of TYK2	PsO
<b>Xeljanz</b> <sup>®</sup> (tofacitinib tablets/oral solution)	Inhibition of JAK pathways	RA, PJIA, PsA, UC
<b>Xeljanz</b> <sup>®</sup> XR (tofacitinib extended-release tablets)	Inhibition of JAK pathways	RA, PsA, UC
<b>Zeposia</b> <sup>®</sup> (ozanimod tablets)	Sphingosine 1 phosphate receptor modulator	UC
<b>Velsipity</b> <sup>®</sup> (etrasimod tablets)	Sphingosine 1 phosphate receptor modulator	UC

\* Not an all-inclusive list of indications. Refer to the prescribing information for the respective agent for FDA-approved indications; SC – Subcutaneous; TNF – Tumor necrosis factor; AS – Ankylosing spondylitis; CD – Crohn’s disease; JIA – Juvenile idiopathic arthritis; PsO – Plaque psoriasis; PsA – Psoriatic arthritis; RA – Rheumatoid arthritis; UC – Ulcerative colitis; nr-axSpA – Non-radiographic axial spondyloarthritis; IV –

Intravenous, PJIA – Polyarticular juvenile idiopathic arthritis; IL – Interleukin; SJIA – Systemic juvenile idiopathic arthritis; ^ Off-label use of Kineret in JIA supported in guidelines; ERA – Enthesitis-related arthritis; DMARD – Disease-modifying antirheumatic drug; PDE4 – Phosphodiesterase 4; JAK – Janus kinase; AD – Atopic dermatitis; AA – Alopecia areata; TYK2 – Tyrosine kinase 2.

## Revision Details

Type of Revision	Summary of Changes	Date
New	New policy	11/1/2024

The policy effective date is in force until updated or retired.

---

“Cigna Companies” refers to operating subsidiaries of The Cigna Group. All products and services are provided exclusively by or through such operating subsidiaries, including Cigna Health and Life Insurance Company, Connecticut General Life Insurance Company, Evernorth Behavioral Health, Inc., Cigna Health Management, Inc., and HMO or service company subsidiaries of The Cigna Group. © 2024 The Cigna Group.