

Fax completed form to: (855) 840-1678
If this is an URGENT request, please call (800) 882-4462 (800.88.CIGNA)

## **Opdivo** (nivolumab)

PHYSICIAN INFORMATION		PATIENT INFORMATION				
* Physician Name:  Specialty: * DEA, NPI or TIN:			*Due to privacy regulations we will not be able to respond via fax with the outcome of our review unless all asterisked (*) items on this form are completed *			
Office Contact Person:			this form are completed.*  * Patient Name:			
Office Phone:			* Cigna ID: * Date of Birth:			
Office Fax:			* Patient Street Address:			
					4	7:
Office Street Address:		T	City: State: Zip:		Zip:	
City:	State:	Zip:	Patient Phone:			
Urgency:  ☐ Standard ☐ Urgent (In checking this box, I attest to the fact that applying the standard review time frame may seriously jeopardize the customer's life, health, or ability to regain maximum function)						
Medication Requested: [	Opdivo 40mg	vial Dpdivo	100mg vial	Opdivo 1	20 mg vial	☐ Opdivo 240mg vial
Directions for use:		Quantity:	Duration of	therapy:	J	-Code:
ICD10:						
Where will this medication be obtained?  ☐ Accredo Specialty Pharmacy** ☐ Prescriber's office stock (billing on a medical claim form) ☐ Other (please specify):  **Cigna's nationally preferred specialty pharmacy  **Medication orders can be placed with Accredo via E-prescribe - Accredo (1620 Century Center Pkwy, Memphis, TN 38134-8822   NCPDP 4436920), Fax 888.302.1028, or Verbal 866.759.1557						
Facility Name:	Facility and/or doctor dispensing and administering medication: Facility Name: State: Tax ID#: Address (City, State, Zip Code):					
Is the patient a candidate for home infusion? Does the physician have an in-office infusion site?						′es □ No □ ′es □ No □
Is the requested medication the patient?	for a chronic or	long-term condition	for which the presci	ription med	lication may be ı	necessary for the life of Yes No
Diagnosis related to use:  ampullary adenocarcinor anal cell carcinoma anaplastic thyroid carcino biliary tract carcinoma bone cancer (including of brain metastases cervical carcinoma chronic lymphocytic leuke colorectal cancer (CRC) endometrial carcinoma esophageal adenocarcin esophageal squamous of esophageal cancer extranodal NK/T-cell lym gastric cancer gastroesophageal junctio gestational trophoblastic hepatocellular carcinoma Hodgkin lymphoma (HL) Kaposi sarcoma	oma hondrosarcoma emia/small lymp oma ell carcinoma (E phoma, nasal ty on (GEJ) cancer neoplasia (GTN	shocytic lymphoma fo SSCC)		·	rmation to diffus	se large B-cell lymphoma

□ malignant pleural mesothelioma (MPM)   □ melanoma   □ Merkel cell carcinoma (MCC)   □ Non-pancreatic neuroendocrine tumor (non-pNET)   □ non-small cell lung cancer (NSCLC)   □ primary mediastinal large B-cell lymphoma (PMLBCL)   □ renal cell carcinoma (RCC)   □ small bowel adenocarcinoma (SBA)   □ soft tissue sarcomas (including angiosarcoma, those of the extremities/body wall/head/neck/retroperitoneal/intra-arhabdomyosarcoma)   □ squamous cell vulvar carcinoma   □ small cell lung cancer (SCLC)   □ urothelial carcinoma (UCC, also transitional cell carcinoma [TCC])   □ squamous cell carcinoma of the head and neck (SCCHN)   □ other (please specify):	abdominal, and	
Clinical Information  Is this new start or continuation of therapy?	gression while on Yes  □ No □	
***This drug requires supportive documentation (i.e. genetic testing, chart notes, lab/test results, etc). Supportive documentation for all answers must be attached with this request.	ortive	
(if anal cell carcinoma, endometrial, non-pNET, squamous vulvar) Was your patient previously treated with only one chemotherapy regimen for this diagnosis?	other	
(if non-pNET and previous first line chemo) Did your patient experience progression (or disease worsening) while on chemotherapy? (if endometrial) Does your patient have recurrent or metastatic disease? (if not recurrent or metastatic) Does your patient have high-risk mismatch repair deficient (dMMR) tumors?	Yes No Series No	
(if bone cancer) Does your patient have tissue tumor mutation burden-high (TMB-H) tumors with 10 or more mutation megabase? (if bone cancer) Has your patient previously been treated with any therapy for this diagnosis?	ns per Yes	
(if yes) Did your patient have disease progression with the previous treatment?	Yes 🗌 No 🗌	
(if bone cancer) Are there any satisfactory alternative options available for treatment?	Yes 🗌 No 🗌	
(if brain mets) Is melanoma the primary tumor/site?	Yes 🗌 No 🗌	
(if no) What is the primary tumor/site (if brain mets) Does your patient have recurrent disease?		
(if CRC) Does your patient have unresectable, advanced, or metastatic disease? (if CRC or SBA) Has your patient undergone immunohistochemistry (IHC) or microsatellite instability (MSI) testing? (if yes) What were the results?  ☐ deficient mismatch repair (dMMR) or microsatellite instability-high (MSI-H)	Yes No No Yes No	
proficient mismatch repair (pMMR) or microsatellite instability-low or stable (MSI-low or MSI-stable)	v	
(if CRC) Has your patient previously used any type of chemotherapy for this diagnosis?	Yes ☐ No ☐	
(if no previous chemo) Is intensive therapy appropriate for your patient?	Yes No No	
(if previous chemo) Has your patient been previously treated with an oxaliplatin-based chemotherapy regimen?	Yes No	
(if CLL/SLL) Does your patient have the del(17p)/TP53 mutation? (if no) Is your patient refractory to chemotherapy and unable to receive chemoimmunotherapy?	Yes  No Yes No No	
(if ESCC) Is this the first therapy your patient has received for this diagnosis?	Yes 🗌 No 🗌	
(if yes) Will your patient be using Opdivo in combination with Yervoy (ipilimumab)?	Yes 🗌 No 🗌	
(if no) Was the patient previously treated with fluoropyrimidine (like capecitabine [Xeloda], floxuridine, and fl Adrucil]) and platinum-based chemotherapy (like carboplatin or cisplatin) for this diagnosis?	uorouracil [5-FU, Yes	
(if yes) Does the patient have unresectable advanced, or metastatic disease?	Yes ☐ No ☐	
(if gastric, GEJ or esophageal adenocarcinoma) Does your patient have advanced or metastatic disease?  (if yes) Is/Will the requested medication (be)ing given in combination with a fluoropyrimidine (like capecitable floxuridine, and fluorouracil [5-FU, Adrucil]) and platinum (like carboplatin or cisplatin)-containing chemother		

	Yes $\square$	No □	
(if esophageal OR GEJ cancer except ESCC) Was your patient treated with chemoradiation followed by surgery to recancer, but some cancer cells were found in the removed tumor or lymph nodes?			
(if yes) Is this medication being given to help prevent the cancer from coming back?	Yes 🗌	No 🗌	
(if extranodal NK/T-cell lymphoma [nasal type] or PMLBCL) Does your patient have relapsed or refractory disease?	Yes 🗌	No 🗌	
(if extranodal NK/T-cell lymphoma, nasal type) Was your patient previously treated with more than 1 regimen of chemical states and the companies of the compani			
(if yes) Was one of the lines of therapy an alternate combination chemotherapy regimen (asparaginase-base previously used?			
(if GTN) Does your patient have recurrent or progressive disease? (if GTN) Was your patient previously treated with a platinum/etoposide-containing regimen? (if HCC) Was your patient previously treated with Nexavar?	Yes 🗌	No 🗌 No 🗍 No 🗍	
(if HL) Which type of Hodgkin lymphoma does your patient have?  classical type nodular lymphocyte predominant type unknown  (if HL) Which of the following applies to your patient? relapsed or refractory disease palliative therapy and patient is older than 60 years neither of the above			
(if relapsed/refractory) Has your patient undergone an autologous stem cell transplant? (if yes) After the transplant, did your patient have therapy with Adcetris?	=	No 🗌 No 🗍	
(if melanoma) How is this medication being used for this diagnosis?  Adjuvant treatment for metastatic disease that has spread to the lymph nodes  Adjuvant treatment for stage IIB/C disease  Single-agent therapy  In combination with ipilimumab (generic for Yervoy)  Other			
(if bone cancer OR melanoma & and not adjuvant) Does your patient have metastatic or unresectable disease?	Yes 🗌	No 🗌	
(if melanoma & adjuvant tx) Did your patient have complete resection of the melanoma?	Yes 🗌	No 🗌	
(if MPM) Which of the following applies?  ☐ Drug requested is being used as single-agent therapy ☐ Drug requested is being given in combination with Yervoy ☐ other			
(if NSCLC) Which best describes Opdivo's role in therapy?  Opdivo is being given as first line treatment Opdivo is being given as subsequent therapy. Opdivo is being given as neoadjuvant therapy. unknown			
(if cervical carcinoma) Is this medication being used as a second line or subsequent therapy?	Yes 🗌	No 🗌	
(if cervical carcinoma) Does the patient have PD-L1 positive disease?	Yes 🗌	No 🗌	
(if bone cancer, non-pNET or NSCLC [1st line]) Is/Will the requested drug be(ing) used in combination with Yervoy (ip	ilimumab) Yes		
(if anal cell carcinoma, non-pNET or NSCLC [1st line or subsequent]) Does your patient have metastatic disease?	Yes 🗌	No 🗌	
(if NSCLC, subsequent) Does your patient have performance status 0-2? (if NSCLC, subsequent) Was your patient previously treated with platinum-base chemotherapy, such as carboplatin control of the cont			
(if not treated before with platinum or NSCLC [1st line]) Which of the following applies to your patient?  ☐ ALK-positive disease ☐ EGFR mutation-positive disease ☐ testing did not indicate either EGFR mutation- or ALK- positive disease ☐ molecular testing was not done	⊔	. •• U	
(if ALK-pos) Was your patient previously treated with Xalkori or Zykadia? (if EGFR mutation-pos) Was your patient previously treated with Gilotrif, Iressa or Tarceva?	_ =	No 🗌 No 🗌	
(if NSCLC, 1st line) Does your patient have PD-L1 expressing (greater than 1%) tumors?	Yes 🗌	No 🗌	

(if NSCLC, neoadjuvant) Does the patient have early-stage disease (stage 1, 2, or 3A)? (if NSCLC, neoadjuvant) Does the patient have resectable disease? (if NSCLC, neoadjuvant) Is/Will the medication be(ing) given with platinum therapy (carboplatin, cisplatin)?	Yes
(if PMLBCL) Which of the following best describes how the requested drug will be given to this patient?  ☐ single agent therapy ☐ given with Adcetris (brentuximab vedotin) ☐ neither of the above/unknown	
(if EGFR mutation-pos) Was your patient previously treated with Gilotrif, Iressa, or Tarceva (erlotinib)?	Yes 🗌 No 🗌
(if SBA) Does your patient have advanced or metastatic disease? (if SBA) Which of the following best describes how the requested drug will be given to this patient?  ☐ as single agent therapy ☐ in combination with Yervoy (ipilimumab) ☐ neither of the above/unknown	Yes  No
(if SCCHN) Was your patient previously treated with platinum-base chemotherapy, such as carboplatin or cisplatin? (if yes) Did your patient have progression of disease afterwards?  Please provide the following details: drug name, date(s) taken and for how long, and what the documented taking each drug.	Yes 🗌 No 🗌
(if RCC) Does your patient have advanced, stage IV, or relapsed disease? (if RCC) Will the drug requested be used in combination with Yervoy? (if yes) Has your patient received any other chemotherapy before for this diagnosis? (if RCC, not in combo with Yervoy) Will the drug requested be used in combination with Cabometyx? (if RCC, with Cabometyx) Is this the first therapy your patient has received for this diagnosis? (if RCC, not in combo with Yervoy or Cabometyx) Has your patient previously received anti-angiogenic therapy (for exavstin, Inlyta, Nexavar, Sutent, Votrient? (if squamous vulvar) Does your patient have HPV-related advanced, recurrent or metastatic disease?	Yes
(if anal cell carcinoma, cervical carcinoma, CRC, endometrial, GTN, HL, NSCLC [not in combo with Yervoy], SCCHN vulvar or RCC) Is the drug requested being used as single-agent therapy?	, squamous cell Yes
(if SCLC) Was your patient previously treated with platinum-based chemotherapy (such as carboplatin or cisplatin) A other line of therapy?  (if yes) Did your patient have progression of disease after these treatments?  Please provide the following details: drug name, date(s) taken and for how long, and what the documented results each drug.	Yes No No Yes No
(if UCC/TCC) Which of these best describes the use of the requested medication?  ☐ As adjuvant treatment in patient at high risk of recurrence after undergoing radical resection ☐ As first line treatment ☐ For locally advanced or metastatic disease ☐ None of the above	
(if UCC/TCC and locally advanced or metastatic) Was your patient previously treated with platinum-base chemothers carboplatin or cisplatin?  (if yes) Did your patient have progression of disease while on the drug or afterwards?  (if UCC/TCC, and first line treatment) Will this medication be used in combination with cisplatin and gemcitabine?  (if UCC/TCC, and first line treatment) Does the patient have metastatic or unresectable disease?	apy, such as Yes  No  Yes
(if biliary tract carcinoma) Has the patient previously been treated with any therapy for this diagnosis? (if biliary tract carcinoma) Has the patient previously been treated with any systemic therapy for this diagnosis? (if biliary tract carcinoma) Is this medication being used as a single agent? (if biliary tract carcinoma) Will your patient be using Opdivo in combination with ipilimumab (generic for Yervoy)? (if biliary tract carcinoma) Was your patient previously treated with a checkpoint inhibitor? (if biliary tract carcinoma) Does your patient have tumor mutation burden-high (TMB-H) tumors? (if biliary tract carcinoma) Did your patient have progression of disease during or after previous systemic treatments? (if biliary tract carcinoma) Does the patient have unresectable, resected gross residual disease, or metastatic disease.	
(if Kaposi sarcoma) Does your patient have relapsed OR refractory disease? (if Kaposi sarcoma) Does your patient have advanced cutaneous, oral, visceral, or nodal disease? (if Kaposi sarcoma) Did your patient have disease progression while on, or did not respond to, first line systemic ther (if Kaposi sarcoma) Will your patient be using Opdivo in combination with ipilimumab (generic for Yervoy)?	Yes ☐ No ☐ Yes ☐ No ☐ apy? Yes ☐ No ☐ Yes ☐ No ☐

(if anaplastic thyroid) Is this medication being used as a single-agent? (if anaplastic thyroid) Does the patient have stage IVC (metastatic) disease?	Yes  No Yes No No
(if anaplastic thyroid) How will this drug be used?  ☐ As aggressive first-line therapy ☐ As second-line therapy ☐ None of the above	
**This drug requires supportive documentation (i.e. genetic testing, chart notes, lab/test results documentation for all answers must be attached with this request.	s, etc). Supportive
<b>Additional Pertinent Information:</b> (including disease stage, prior therapy, performance status, and not any agents to be used concurrently).	ames/doses/admin schedule of
Attestation: I attest the information provided is true and accurate to the best of my knowledge. I under insurer its designees may perform a routine audit and request the medical information necessary information reported on this form.	
Prescriber Signature: Date:	
Save Time! Submit Online at: www.covermymeds.com/main/prior-authorization-forms/cigna/	r via SureScripts in your EHR.
Our standard response time for prescription drug coverage requests is 5 business days. If your request you call us to expedite the request. View our Prescription Drug List and Coverage Policies	

v081524

"Cigna" is a registered service mark, and the "Tree of Life" logo is a service mark, of Cigna Intellectual Property, Inc., licensed for use by Cigna Corporation and its operating subsidiaries. All products and services are provided by or through such operating subsidiaries and not by Cigna Corporation. Such operating subsidiaries include, for example, Cigna Health and Life Insurance Company and Cigna Health Management, Inc. Address: Cigna Pharmacy Services, PO Box 42005, Phoenix AZ 85080-2005